

Department of Mathematics and Statistics


Preprint MPS-2016-03

15 March 2016

A Sub-Riemannian Santalo Formula with Applications to Isoperimetric Inequalities and First Dirichlet Eigenvalue of Hypoelliptic Operators

by

Dario Prandi, Luca Rizzi and **Marcello Seri**



A SUB-RIEMANNIAN SANTALÓ FORMULA WITH APPLICATIONS TO ISOPERIMETRIC INEQUALITIES AND FIRST DIRICHLET EIGENVALUE OF HYPOELLIPTIC OPERATORS

DARIO PRANDI¹, LUCA RIZZI², AND MARCELLO SERI³

ABSTRACT. Sub-Riemannian geometry is a generalization of Riemannian one, to include nonholonomic constraints. In this paper we prove a nonholonomic version of the classical Santaló formula: a result in integral geometry that describes the intrinsic Liouville measure on the unit cotangent bundle in terms of the geodesic flow. Our construction works under quite general assumptions, satisfied by any sub-Riemannian structure associated with a Riemannian foliation with totally geodesic leaves (e.g. CR and quaternionic contact manifolds with symmetries) and any Carnot group. A key ingredient is a “reduction procedure” that allows to consider only a simple subset of sub-Riemannian geodesics.

As an application, we derive (p-)Hardy-type and isoperimetric-type inequalities for a compact domain M with piecewise C^2 boundary. Moreover, we prove a universal (i.e. curvature independent) lower bound for the first Dirichlet eigenvalue $\lambda_1(M)$ of the intrinsic sub-Laplacian,

$$\lambda_1(M) \geq \frac{k\pi^2}{L^2},$$

in terms of the rank k of the distribution and the length L of the longest reduced sub-Riemannian geodesic contained in M . All our results are sharp for the sub-Riemannian structures on the hemispheres of the complex and quaternionic Hopf fibrations:

$$\mathbb{S}^1 \hookrightarrow \mathbb{S}^{2d+1} \xrightarrow{p} \mathbb{C}\mathbb{P}^d, \quad \mathbb{S}^3 \hookrightarrow \mathbb{S}^{4d+3} \xrightarrow{p} \mathbb{H}\mathbb{P}^d, \quad d \geq 1,$$

where the sub-Laplacian is the standard hypoelliptic operator of CR and quaternionic contact geometries, $L = \pi$ and $k = 2d$ or $4d$, respectively.

1. INTRODUCTION AND RESULTS

Let (M, g) be a compact Riemannian manifold with boundary ∂M . Santaló formula [19, 43] is a classical result in integral geometry that describes the Liouville measure μ of the unit tangent bundle UM in terms of the geodesic flow $\phi_t : UM \rightarrow UM$. Namely, for any measurable function $F : UM \rightarrow \mathbb{R}$ we have

$$(1) \quad \int_{U^{\circlearrowleft} M} F \mu = \int_{\partial M} \left[\int_{U_q^+ \partial M} \left(\int_0^{\ell(v)} F(\phi_t(v)) dt \right) g(v, \mathbf{n}_q) \eta_q(v) \right] \sigma(q),$$

where σ is the surface form on ∂M induced by the inward pointing normal vector \mathbf{n} , η_q is the Riemannian spherical measure on $U_q M$, $U_q^+ \partial M$ is the set of inward pointing unit vectors at $q \in \partial M$ and $\ell(v)$ is the exit length of the geodesic with initial vector v . Finally,

¹CEREMADE, UNIVERSITÉ PARIS-DAUPHINE, PARIS, FRANCE.

²CNRS, CMAP ÉCOLE POLYTECHNIQUE, AND ÉQUIPE INRIA GECO SACLAY ÎLE-DE-FRANCE, PARIS, FRANCE

³DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF READING, READING, UK
E-mail addresses: prandi@ceremade.dauphine.fr, luca.rizzi@cmap.polytechnique.fr, m.seri@ucl.ac.uk.

2010 *Mathematics Subject Classification.* 53C17, 53C65, 35P15, 57R30, 35R03, 53C65.

This research has been supported by the European Research Council, ERC StG 2009 “GeCoMethods”, contract n. 239748, by the iCODE institute (research project of the Idex Paris-Saclay), by the SMAI project “BOUM”. The first and second authors were partially supported by the Grant ANR-15-CE40-0018 of the ANR. This research benefited from the support of the “FMJH Program Gaspard Monge in optimization and operation research” and from the support to this program from EDF..

$U^*M \subseteq UM$ is the visible set, i.e. the set of unit vectors that can be reached via the geodesic flow starting from points on ∂M .

In the Riemannian setting, (1) allows to deduce some very general and non-trivial isoperimetric inequalities and Dirichlet eigenvalues estimates for the Laplace-Beltrami operator as showed by Croke in the celebrated papers [21, 22, 23].

The extension of (1) to the sub-Riemannian setting and its consequences are not straightforward for a number of reasons. Firstly, in sub-Riemannian geometry the geodesic flow is replaced by a degenerate Hamiltonian flow on the cotangent bundle. Moreover, the unit cotangent bundle (the set of covectors with unit norm) is not compact, but rather has the topology of an infinite cylinder. Finally, in sub-Riemannian geometry there is not a clear agreement on which is the “canonical” volume, generalizing the Riemannian measure. Another aspect to consider is the presence of characteristic points on the boundary.

In this paper we extend (1) to the most general class of sub-Riemannian structures for which Santaló formula makes sense. As an application we deduce Hardy-like inequalities, sharp universal estimates on the first Dirichlet eigenvalue of the sub-Laplacian and sharp isoperimetric-type inequalities.

To our best knowledge, a sub-Riemannian version of (1) appeared only in [40] for the three-dimensional Heisenberg group, and more recently in [38] for Carnot groups, where the natural global coordinates allow for explicit computations. As far as other sub-Riemannian structures are concerned, Santaló formula is an unexplored technique with potential applications to different settings, including CR (Cauchy-Riemann) and QC (quaternionic contact) geometry, Riemannian foliations, and Carnot groups.

1.1. Setting and examples. Let (N, \mathcal{D}, g) be a sub-Riemannian manifold of dimension n , where $\mathcal{D} \subseteq TN$ is a distribution that satisfies the bracket-generating condition and g is a smooth metric on \mathcal{D} . Sections $X \in \Gamma(\mathcal{D})$ are called *horizontal*. We consider a compact n -dimensional submanifold $M \subset N$ with boundary $\partial M \neq \emptyset$.

If (N, \mathcal{D}, g) is Riemannian, we equip it with its Riemannian volume ω_R . In the genuinely sub-Riemannian case we fix any smooth volume form ω on M (or a density if M is not orientable). In any case, the surface measure $\sigma = \iota_{\mathbf{n}}\omega$ on ∂M is given by the contraction with the horizontal unit normal \mathbf{n} to ∂M . For what concerns the regularity of the boundary, we assume only that ∂M is piecewise C^2 . (See also Remark 6 for a discussion of the case of Lipschitz boundary.)

A central role is played by sub-Riemannian geodesics, i.e., curves tangent to \mathcal{D} that locally minimize the sub-Riemannian distance between endpoints. In this setting, the geodesic flow¹ is a natural Hamiltonian flow $\phi_t : T^*M \rightarrow T^*M$ on the cotangent bundle, induced by the Hamiltonian function $H \in C^\infty(T^*M)$. The latter is a non-negative, degenerate, quadratic form on the fibers of T^*M that contains all the information on the sub-Riemannian structure. Length-parametrized geodesics are characterized by an initial covector λ in the unit cotangent bundle $U^*M = \{\lambda \in T^*M \mid 2H(\lambda) = 1\}$.

A key ingredient for most of our results is the following *reduction procedure*. Fix a transverse sub-bundle $\mathcal{V} \subset TM$ such that $TM = \mathcal{D} \oplus \mathcal{V}$. We define the reduced cotangent bundle T^*M^r as the set of covectors annihilating \mathcal{V} . On T^*M^r we define a reduced Liouville volume Θ^r , which depends on the choice of the volume ω on M . These must satisfy the following stability hypotheses:

- (H1) The bundle T^*M^r is invariant under the Hamiltonian flow ϕ_t ;
- (H2) The reduced Liouville volume is invariant, i.e. $\mathcal{L}_{\bar{H}}\Theta^r = 0$.

This allows to reduce the non-compact U^*M to a compact slice $U^*M^r := U^*M \cap T^*M^r$, equipped with an invariant measure (see Section 4.3). These hypotheses are verified for:

¹Abnormal geodesics are allowed, but strictly abnormal ones, not given by the Hamiltonian flow on the cotangent bundle, do not play any role in our construction.




FIGURE 1. Exit length (left) and visible set (right). Covectors are represented as hyperplanes, the arrow shows the direction of propagation of the associated geodesic for positive time.

- any Riemannian structure, equipped with the Riemannian volume;
- any sub-Riemannian structure associated with a Riemannian foliation with totally geodesic leaves, equipped with the Riemannian volume (including contact, CR, QC structures with transverse symmetries), see Section 5.2;
- any left-invariant sub-Riemannian structure on a Carnot group², equipped with the Haar volume, see Section 5.1.

An interesting example, coming from CR geometry, is the complex Hopf fibration (CHF)

$$\mathbb{S}^1 \hookrightarrow \mathbb{S}^{2d+1} \xrightarrow{p} \mathbb{C}\mathbb{P}^d, \quad d \geq 1,$$

where $\mathcal{D} := (\ker p_*)^\perp$ is the orthogonal complement of the kernel of the differential of the Hopf map w.r.t. the round metric on \mathbb{S}^{2d+1} , and the sub-Riemannian metric g is the restriction to \mathcal{D} of the round one. Another interesting structure, coming from QC geometry and with corank 3, is the quaternionic Hopf fibration (QHF)

$$\mathbb{S}^3 \hookrightarrow \mathbb{S}^{4d+3} \xrightarrow{p} \mathbb{H}\mathbb{P}^d, \quad d \geq 1,$$

where $\mathbb{H}\mathbb{P}^d$ is the quaternionic projective space of real dimension $4d$ and the sub-Riemannian structure on \mathbb{S}^{4d+3} is defined similarly to its complex version.

1.2. Sub-Riemannian Santaló formulas. Consider a sub-Riemannian geodesic $\gamma(t)$ with initial covector $\lambda \in U^*M$. The *exit length* $\ell(\lambda) \in [0, +\infty)$ is the length after which γ leaves M by crossing ∂M . Similarly, $\tilde{\ell}(\lambda)$ is the minimum between $\ell(\lambda)$ and the cut length $c(\lambda)$. That is, after length $\tilde{\ell}(\lambda)$ the geodesic either loses optimality or leaves M .

The *visible unit cotangent bundle* $U^{\leftarrow}M \subset U^*M$ is the set of unit covectors λ such that $\ell(-\lambda) < +\infty$. (See Fig. 1.) Analogously, the *optimally visible unit cotangent bundle* $\tilde{U}^{\leftarrow}M$ is the set of unit covectors such that $\tilde{\ell}(-\lambda) < +\infty$.

For any non-characteristic point $q \in \partial M$, we have a well defined inner pointing unit horizontal vector $\mathbf{n}_q \in \mathcal{D}_q$, and $U_q^+ \partial M \subset U_q^*M$ is the set of initial covectors of geodesics that, for positive time, are directed toward the interior of M .

As anticipated, we do not consider all the length-parametrized geodesics, i.e. all initial covectors $\lambda \in U_q^*M \simeq \mathbb{S}^{k-1} \times \mathbb{R}^{n-k}$, but a *reduced subset* $U_q^*M^r \simeq \mathbb{S}^{k-1}$. In the following the suffix r always denotes the intersection with the reduced unit cotangent bundle U^*M^r . We stress the critical fact that U^*M^r is compact, while U^*M never is, except in the

²We stress that Carnot groups are not Riemannian foliations if their step is > 2 .

Riemannian setting where the reduction procedure is trivial. With these basic definitions at hand, we are ready to state the sub-Riemannian Santaló formulas.

Theorem 1 (Reduced Santaló formulas). *The visible set $U^{\diamond}M^r$ and the optimally visible set $\tilde{U}^{\diamond}M^r$ are measurable. For any measurable function $F : U^*M^r \rightarrow \mathbb{R}$ we have*

$$(2) \quad \int_{U^{\diamond}M^r} F \mu^r = \int_{\partial M} \left[\int_{U_q^+ \partial M^r} \left(\int_0^{\ell(\lambda)} F(\phi_t(\lambda)) dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q^r(\lambda) \right] \sigma(q),$$

$$(3) \quad \int_{\tilde{U}^{\diamond}M^r} F \mu^r = \int_{\partial M} \left[\int_{U_q^+ \partial M^r} \left(\int_0^{\tilde{\ell}(\lambda)} F(\phi_t(\lambda)) dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q^r(\lambda) \right] \sigma(q).$$

In (2)-(3), μ^r is a reduced invariant Liouville measure on U^*M^r , η_q^r is an appropriate smooth measure on the fibers $U_q^*M^r$ and $\langle \lambda, \cdot \rangle$ denotes the action of covectors on vectors. Indeed both include the Riemannian case, where the reduction procedure is trivial and $U^*M \simeq UM$ since the Hamiltonian is not degenerate.

Remark 1. Hypotheses **(H1)** and **(H2)** are essential for the reduction procedure. An unreduced version of Theorem 1 holds for any volume ω and with no other assumptions but the Lipschitz regularity of ∂M (see Theorem 16 and Remark 6). However, the consequences we present do not hold a priori, as their proofs rely on the summability of certain functions on U^*M^r , generally false on the non-compact U^*M .

1.3. Hardy-type inequalities. For any $f \in C^\infty(M)$, let $\nabla_H f \in \Gamma(\mathcal{D})$ be the *horizontal gradient*: the horizontal direction of steepest increase of f . It is defined via the identity

$$g(\nabla_H f, X) = df(X), \quad \forall X \in \Gamma(\mathcal{D}).$$

Consider all length-parametrized sub-Riemannian geodesic passing through a point $q \in M$, with covector $\lambda \in U_q^*M$. Set $L(\lambda) := \ell(\lambda) + \ell(-\lambda)$; this is the length of the maximal geodesic that passes through q with covector λ .

Proposition 2 (Hardy-like inequalities). *For any $f \in C_0^\infty(M)$ it holds*

$$(4) \quad \int_M |\nabla_H f|^2 \omega \geq \frac{k\pi^2}{|\mathbb{S}^{k-1}|} \int_M \frac{f^2}{R^2} \omega,$$

$$(5) \quad \int_M |\nabla_H f|^2 \omega \geq \frac{k}{4|\mathbb{S}^{k-1}|} \int_M \frac{f^2}{r^2} \omega,$$

where $k = \text{rank } \mathcal{D}$ and $r, R : M \rightarrow \mathbb{R}$ are:

$$\frac{1}{R^2(q)} := \int_{U_q^*M^r} \frac{1}{L^2} \eta_q^r, \quad \frac{1}{r^2(q)} := \int_{U_q^*M^r} \frac{1}{\ell^2} \eta_q^r, \quad \forall q \in M.$$

We observe that r is the *harmonic mean distance from the boundary* defined in [25]. One can also consider the following generalization of Proposition 2 for $L^p(M, \omega)$ norms.

Proposition 3 (p -Hardy-like inequality). *Let $p > 1$ and $f \in C_0^\infty(M)$. Then*

$$(6) \quad \int_M |\nabla_H f|^p \omega \geq \pi_p^p C_{p,k} \int_M \frac{|f|^p}{R^p} \omega,$$

$$(7) \quad \int_M |\nabla_H f|^p \omega \geq \left(\frac{p-1}{p} \right)^p C_{p,k} \int_M \frac{|f|^p}{r^p} \omega,$$

where $k = \text{rank } \mathcal{D}$, the constants π_p and $C_{p,k}$ are

$$\pi_p = \frac{2\pi(p-1)^{1/p}}{p \sin(\pi/p)}, \quad C_{p,k} = \frac{k}{|\mathbb{S}^{k-1}|} \frac{2\Gamma(1 + \frac{k}{2})\Gamma(1 + \frac{p}{2})}{\sqrt{\pi}\Gamma(\frac{k+p}{2})},$$

and $r^p, R^p : M \rightarrow \mathbb{R}$ are

$$\frac{1}{R^p(q)} := \int_{U_q^* M^r} \frac{1}{L^p} \eta_q^r, \quad \frac{1}{r^p(q)} := \int_{U_q^* M^r} \frac{1}{\ell^p} \eta_q^r, \quad \forall q \in M.$$

1.4. Lower bound for the first Dirichlet eigenvalue. For any given smooth volume ω , a fundamental operator in sub-Riemannian geometry is the sub-Laplacian Δ_ω , playing the role of the Laplace-Beltrami operator in Riemannian geometry. Under the bracket-generating condition, this is an hypoelliptic operator, self-adjoint on $L^2(M, \omega)$. Its principal symbol is (twice) the Hamiltonian, thus the Dirichlet spectrum of $-\Delta_\omega$ on the compact manifold M is positive and discrete. We denote it

$$0 < \lambda_1(M) \leq \lambda_2(M) \leq \dots$$

As a consequence of Proposition 2 and the min-max principle, we obtain a universal lower bound for the first Dirichlet eigenvalue $\lambda_1(M)$ on the given domain. Here with universal we mean an estimate not requiring any *a priori* assumption on curvature or capacity.

Proposition 4 (Universal spectral lower bound). *Let $L = \sup_{\lambda \in U^* M^r} L(\lambda)$ be the length of the longest reduced geodesic contained in M . Then, letting $k = \text{rank } \mathcal{D}$,*

$$(8) \quad \lambda_1(M) \geq \frac{k\pi^2}{L^2}.$$

In the Riemannian case, as noted by Croke, we attain equality in (8) when M is the hemisphere of the Riemannian round sphere. We prove the following extension to the sub-Riemannian setting.

Proposition 5 (Sharpness of the eigenvalue lower bound). *In Proposition 4, in the following cases we have equality, for all $d \geq 1$:*

- (i) *the hemispheres \mathbb{S}_+^d of the Riemannian round sphere \mathbb{S}^d ;*
- (ii) *the hemispheres \mathbb{S}_+^{2d+1} of the sub-Riemannian complex Hopf fibration \mathbb{S}^{2d+1} ;*
- (iii) *the hemispheres \mathbb{S}_+^{4d+3} of the sub-Riemannian quaternionic Hopf fibration \mathbb{S}^{4d+3} ,*
all equipped with the Riemannian volume of the corresponding round sphere. In all these cases, $L = \pi$ and $\lambda_1(M) = d, 2d$ or $4d$, respectively. Moreover, the associated eigenfunction is $\Psi = \cos(\delta)$, where δ is the Riemannian distance from the north pole.

Remark 2. The Riemannian volume of the sub-Riemannian Hopf fibrations coincides, up to a constant factor, with their Popp volume [4, 39], an intrinsic smooth measure in sub-Riemannian geometry [12, Sec. 3]. This is proved for 3-Sasakian structures (including the QHF) in [42, Prop. 34] and can be proved exactly in the same way for Sasakian structures (including the CHF) using the explicit formula for Popp volume of [4]. For the case (i) Δ_ω is the Laplace-Beltrami operator. For the cases (ii) and (iii) Δ_ω is the standard sub-Laplacian of CR and QC geometry, respectively.

In (8), L cannot be replaced by the sub-Riemannian diameter, as M might contain very long (non-optimal) geodesics, for example closed ones, and $L = +\infty$. In principle, L can be computed when the reduced geodesic flow is explicit. This is the case for Carnot groups, where reduced geodesics passing through the origin are simply straight lines (they fill a k -plane for rank k Carnot groups). It turns out that, in this case, $L = \text{diam}_H(M)$ (the *horizontal diameter*, that is the diameter of the set M measured through left-translations of the aforementioned straight lines). Thus (8) gives an easily computable lower bound for the first Dirichlet eigenvalue in terms of purely metric quantities.

Corollary 6. *Let M be a compact n -dimensional submanifold with piecewise C^2 boundary of a Carnot group of rank k , with the Haar volume. Then,*

$$(9) \quad \lambda_1(M) \geq \frac{k\pi^2}{\text{diam}_H(M)^2},$$




FIGURE 2. Visibility angle on a 2D Riemannian manifold. Only the geodesics with tangent vector in the dashed slice go to ∂M .

where diam_H denotes the horizontal diameter of the Carnot group.

In particular, if M is the metric ball of radius R , we obtain $\lambda_1(M) \geq k\pi^2/(2R)^2$. Clearly (9) is not sharp, as one can check easily in the Euclidean case.

1.5. Isoperimetric-type inequalities. In this section we relate the sub-Riemannian area and perimeter of M with some of its geometric properties. Since M is compact, the sub-Riemannian diameter $\text{diam}(M)$ can be characterized as the length of the longest optimal geodesic contained in M . Analogously, the reduced sub-Riemannian diameter $\text{diam}^r(M)$ is the length of the longest reduced optimal geodesic contained in M . Indeed $\text{diam}^r(M) \leq \text{diam}(M)$.

Consider all (reduced) geodesics passing through $q \in M$ with covector λ . Some of them originate from the boundary ∂M , that is $\ell(-\lambda) < +\infty$; others do not, i.e. $\ell(-\lambda) = +\infty$. The relative ratio of these two types of geodesics (w.r.t. an appropriate measure on $U_q^*M^r$) is called the *visibility angle* $\vartheta_q^\star \in [0, 1]$ at q (see Definition 6). Roughly speaking, if $\vartheta_q^\star = 1$ then any geodesic passing through q will hit the boundary and, on the opposite, if it is equal to 0 then q is not visible from the boundary (see Fig. 2). Similarly, we define the *optimal visibility angle* $\tilde{\vartheta}_q^\star$ by replacing $\ell(-\lambda)$ with $\tilde{\ell}(-\lambda)$. Finally, the *least visibility angle* is $\vartheta^\star := \inf_{q \in M} \vartheta_q^\star$, and similarly for the *least optimal visibility angle* $\tilde{\vartheta}^\star := \inf_{q \in M} \tilde{\vartheta}_q^\star$.

Proposition 7 (Isoperimetric-type inequalities). *Let $\ell := \sup\{\ell(\lambda) \mid \lambda \in U_q^*M^r, q \in \partial M\}$ be the length of the longest reduced geodesic contained in M starting from the boundary ∂M . Then*

$$(10) \quad \frac{\sigma(\partial M)}{\omega(M)} \geq C \frac{\vartheta^\star}{\ell} \quad \text{and} \quad \frac{\sigma(\partial M)}{\omega(M)} \geq C \frac{\tilde{\vartheta}^\star}{\text{diam}^r(M)},$$

where $C = 2\pi|\mathbb{S}^{k-1}|/|\mathbb{S}^k|$ and we set the r.h.s. to 0 if $\ell = +\infty$.

The equality in (10) holds for the hemisphere of the Riemannian round sphere, as pointed out in [21]. We have the following generalization to the sub-Riemannian setting.

Proposition 8 (Sharpness of isoperimetric inequalities). *In Proposition 7, in the following cases we have equality, for all $d \geq 1$:*

- (i) the hemispheres \mathbb{S}_+^d of the Riemannian round sphere \mathbb{S}^d ;
- (ii) the hemispheres \mathbb{S}_+^{2d+1} of the sub-Riemannian complex Hopf fibration \mathbb{S}^{2d+1} ;
- (iii) the hemispheres \mathbb{S}_+^{4d+3} of the sub-Riemannian quaternionic Hopf fibration \mathbb{S}^{4d+3} ;

where ω is the Riemannian volume of the corresponding round sphere. In all these cases $\vartheta^\star = \tilde{\vartheta}^\star = 1$ and $\ell = \text{diam}^r(M) = \pi$.

We can apply Proposition 7 to Carnot groups equipped with the Haar measure. In this case $\vartheta^{\check{\bullet}} = \tilde{\vartheta}^{\check{\bullet}} = 1$ and $\ell = \text{diam}^r(M) = \text{diam}_H(M)$. Moreover, ω is the Lebesgue volume of \mathbb{R}^n and σ is the associated perimeter measure of geometric measure theory [16].

Corollary 9. *Let M be a compact n -dimensional submanifold with piecewise C^2 boundary of a Carnot group of rank k , with the Haar volume. Then,*

$$\frac{\sigma(\partial M)}{\omega(M)} \geq \frac{2\pi|\mathbb{S}^{k-1}|}{|\mathbb{S}^k| \text{diam}_H(M)},$$

where $\text{diam}_H(M)$ is the horizontal diameter of the Carnot group.

This inequality is not sharp even in the Euclidean case, but it is very easy to compute the horizontal diameter for explicit domains. For example, if M is the sub-Riemannian metric ball of radius R , then $\text{diam}_H(M) = 2R$.

1.6. Remark on the change of volume. Fix a sub-Riemannian structure (N, \mathcal{D}, g) , a compact set M with piecewise C^2 boundary and a complement \mathcal{V} such that **(H1)** holds. Now assume that, for some choice of volume form ω , also **(H2)** is satisfied, so that we can carry on with the reduction procedure and all our results hold. One can derive the analogous of Propositions 2, 3, 4, 7 for any other volume $\omega' = e^\varphi \omega$, with $\varphi \in C^\infty(M)$. In all these results, it is sufficient to multiply the r.h.s. of the inequalities by the *volumetric constant* $0 < \alpha \leq 1$ defined as $\alpha := \frac{\min e^\varphi}{\max e^\varphi}$, and indeed replace ω with $\omega' = e^\varphi \omega$ in Propositions 2 and 3, σ with $\sigma' = e^\varphi \sigma$ in Proposition 7, and the sub-Laplacian Δ_ω with³ $\Delta_{\omega'}$ in Proposition 4. Analogously, one can deal with the Corollaries 6 and 9 about Carnot groups.

This remark allows, for example, to obtain results for (sub-)Riemannian weighted measures. This is particularly interesting in the genuinely sub-Riemannian setting since, in some cases, the volume satisfying **(H2)** might not coincide with the intrinsic Popp one.

1.7. Afterwords and further developments. Despite its broad range of applications in Riemannian geometry and its Finsler generalizations [45], only a few works used Santaló formula in the hypoelliptic setting, all of them in the specific case of Carnot groups [38, 40] or 3D Sasakian structures [17]. It is interesting to notice that, in [40], Pansu was able to use it in pairs with minimal surfaces to eliminate the diameter term in Corollary 9 and obtain his celebrated isoperimetric inequality. In our general setting, this is something worth investigating.

The study of spectral properties of hypoelliptic operators is an active area of research. Many results are available for the complete spectrum of the sub-Laplacian on *closed* manifolds (with no boundary conditions). We recall [6, 8, 9] for the case of SU(2), CHF and QHF. Furthermore, in [20], one can find the spectrum of the “flat Heisenberg case” (a compact quotient of the Heisenberg group) together with quantum ergodicity results for 3D contact sub-Riemannian structures.

Concerning the Dirichlet spectrum, the problem is complicated by the presence of the boundary. An account of the existing results in the Riemannian setting, starting from those of Li-Yau and Yang-Zhong, can be found in [18]. In the sub-Riemannian setting, we are aware of the results for the sum of Dirichlet eigenvalues [44] by Strichartz and related spectral inequalities [30] by Hannson and Laptev, both for the case of the Heisenberg group. To our best knowledge, Proposition 4 is the first sharp universal lower bound for the first Dirichlet eigenvalue in the hypoelliptic setting and in particular for non-Carnot structures. Most certainly our sharpness result holds also for the hemisphere of the exotic sub-Riemannian structure on the octonionic Hopf fibration (OHF) $\mathbb{S}^7 \hookrightarrow \mathbb{S}^{15} \rightarrow \mathbb{O}\mathbb{P}^1$ whose spectral properties have never been studied from the sub-Riemannian viewpoint.

³Indeed $\Delta_{\omega'} u = \Delta_\omega u + \langle d\varphi, \nabla_H u \rangle$ for all $u \in C^\infty(M)$.

We observe that the CHF, QHF and OHF are the only Riemannian submersions of the sphere with totally geodesic fibers [27].

The study of Hardy's inequalities, already in the Euclidean setting, ranges across the last century and continues to the present day (see [3, 14, 26] and references therein). The sub-Riemannian case is more recent, for an account of the known result we mention the works for Carnot groups of Capogna, Danielli and Garofalo (see e.g. [15, 24]).

Poincaré inequalities are strictly connected to Hardy's ones. On this subject the literature is again huge, we already mentioned the works of Croke and Derdzinski concerning the Riemannian case [21, 22, 23]. Finally, there is a detailed survey by Ivanov and Vassiliev [33] reviewing the results for CR and QC manifold under Ricci curvature assumptions in the spirit of the Lichnerowicz-Obata theorem. Lower bounds for the first (non-zero) eigenvalue of the sub-Laplacian on closed foliation, under curvature-like assumptions, appeared in [7] (see also [5] for a more general statement).

Possible developments that deserve further investigation are rigidity results in the spirit of Croke [21, 23]. In fact, it seems reasonable that the CHF or QHF are essentially the only sub-Riemannian structures on foliations with totally geodesic leaves (with corank 1 or 3) such that equality holds in Proposition 4 or 7.

We mention also that Santaló formula has possible applications to the problem of self-adjointness of Laplace-Beltrami operators on almost-Riemannian structures [11, 13]. Finally, our Hamiltonian approach is flexible enough to address a very broad field of applications to geometries whose geodesic flow is a Hamiltonian one and lives on stable sub-bundle of the cotangent bundle.

In this paper we focused mostly on foliations, where our results are sharp. For Carnot groups, Corollaries 6 and 9 appeared in [38] and are not sharp. Let us consider for simplicity the 3D Heisenberg group, with coordinates $(x, y, z) \in \mathbb{R}^3$. A relevant class of domains for the Dirichlet eigenvalues problem are the "Heisenberg cubes" $[0, \varepsilon] \times [0, \varepsilon] \times [0, \varepsilon^2]$, obtained by non-homogeneous dilation of the unit cube $[0, 1]^3$. These represent a fundamental domain for the quotient $\mathbb{H}_3/\varepsilon\Gamma$ of the 3D Heisenberg group \mathbb{H}_3 by the (dilation of the) integer Heisenberg subgroup Γ (a lattice). This is the basic example of *nilmanifold*, considered by Gromov in [29], and increasingly being seen as having a role in arithmetic combinatorics and higher-order Fourier analysis [28] (we thank R. Montgomery for pointing out this connection). For these fundamental domains, the first Dirichlet eigenvalue is presently unknown. However, we mention that for any Carnot group the reduction technique developed here can be further improved leading to a λ_1 estimate for cubes, which is sharp in the Euclidean case and, according to numerical experiments, we suspect to be sharp also in \mathbb{H}_3 . This deserves further investigation and will be the subject of a forthcoming paper.

2. SUB-RIEMANNIAN GEOMETRY

We give here only the essential ingredients for our analysis; for more details see [2, 39, 41]. A sub-Riemannian manifold is a triple (M, \mathcal{D}, g) , where M is a smooth, connected manifold of dimension $n \geq 3$, \mathcal{D} is a vector distribution of constant rank $k \leq n$ and g is a smooth metric on \mathcal{D} . We assume that the distribution is bracket-generating, that is

$$(11) \quad \text{span}\{[X_{i_1}, [X_{i_2}, [\dots, [X_{i_{m-1}}, X_{i_m}]]]] \mid m \geq 1\}_q = T_q M, \quad \forall q \in M,$$

for some (and thus any) set $X_1, \dots, X_k \in \Gamma(\mathcal{D})$ of local generators for \mathcal{D} .

A *horizontal curve* $\gamma : [0, T] \rightarrow \mathbb{R}$ is a Lipschitz continuous path such that $\dot{\gamma}(t) \in \mathcal{D}_{\gamma(t)}$ for almost any t . Horizontal curves have a well defined length

$$\ell(\gamma) = \int_0^T \sqrt{g(\dot{\gamma}(t), \dot{\gamma}(t))} dt.$$

Furthermore, the *sub-Riemannian distance* is defined by:

$$d(x, y) = \inf\{\ell(\gamma) \mid \gamma(0) = x, \gamma(T) = y, \gamma \text{ horizontal}\}.$$

By the Chow-Rashevskii theorem, under the bracket-generating condition, d is finite and continuous. Sub-Riemannian geometries include the Riemannian one, when $\mathcal{D} = TM$.

2.1. Sub-Riemannian geodesic flow. *Sub-Riemannian geodesics* are horizontal curves that locally minimize the length between their endpoints. Let $\pi : T^*M \rightarrow M$ be the cotangent bundle. The *sub-Riemannian Hamiltonian* $H : T^*M \rightarrow \mathbb{R}$ is

$$H(\lambda) := \frac{1}{2} \sum_{i=1}^k \langle \lambda, X_i \rangle^2, \quad \lambda \in T^*M,$$

where $X_1, \dots, X_k \in \Gamma(\mathcal{D})$ is any local orthonormal frame and $\langle \lambda, \cdot \rangle$ denotes the action of covectors on vectors. Let σ be the canonical symplectic 2-form on T^*M . The *Hamiltonian vector field* \vec{H} is defined by $\sigma(\cdot, \vec{H}) = dH$. Then the Hamilton equations are

$$(12) \quad \dot{\lambda}(t) = \vec{H}(\lambda(t)).$$

Solutions of (12) are called *extremals*, and their projections $\gamma(t) := \pi(\lambda(t))$ on M are smooth geodesics. The *sub-Riemannian geodesic flow* $\phi_t \in T^*M \rightarrow T^*M$ is the flow of \vec{H} . Thus, any initial covector $\lambda \in T^*M$ is associated with a geodesic $\gamma_\lambda(t) = \pi \circ \phi_t(\lambda)$, and its speed $\|\dot{\gamma}(t)\| = 2H(\lambda)$ is constant. The *unit cotangent bundle* is

$$U^*M = \{\lambda \in T^*M \mid 2H(\lambda) = 1\}.$$

It is a fiber bundle with fiber $U_q^*M = \mathbb{S}^{k-1} \times \mathbb{R}^{n-k}$. For $\lambda \in U_q^*M$, the curve $\gamma_\lambda(t)$ is a *length-parametrized* geodesic with length $\ell(\gamma|_{[t_1, t_2]}) = t_2 - t_1$.

Remark 3. There is also another poorly understood class of locally length minimizing curves, called *abnormal geodesics*, that might not follow the Hamiltonian dynamic of (12). Abnormal geodesics do not exist in Riemannian geometry, and they are all trivial curves in some basic but popular classes of sub-Riemannian structures (e.g. fat ones). Our construction takes in account only the (normal) sub-Riemannian geodesic flow, hence abnormal geodesics are allowed, but ignored. Some hard open problems in sub-Riemannian geometry are related to abnormal geodesics [1, 39] as, for example, the “size” of the set of points reached by abnormal geodesics (see [34, 35] for recent developments).

2.2. The intrinsic sub-Laplacian. Let (M, \mathcal{D}, g) be a compact sub-Riemannian manifold with piecewise C^2 boundary ∂M , and $\omega \in \Lambda^n M$ be any smooth volume form (or a density, if M is not orientable). We define the *Dirichlet energy functional* as

$$E(f) = \int_M 2H(df) \omega, \quad f \in C_0^\infty(M).$$

The Dirichlet energy functional induces the operator $-\Delta_\omega$ on $L^2(M, \omega)$. Its Friedrichs extension is a non-negative self-adjoint operator on $L^2(M, \omega)$ that we call the *Dirichlet sub-Laplacian*. Its domain is the space $H_0^1(M)$, the closure in the $H^1(M)$ norm of the space $C_0^\infty(M)$ of smooth functions that vanish on ∂M . We stress that the sub-Laplacian depends both on the sub-Riemannian structure and on the choice of the volume ω . The spectrum of $-\Delta_\omega$ is discrete and positive,

$$0 < \lambda_1(M) \leq \lambda_2(M) \leq \dots \rightarrow +\infty.$$

In particular, by the min-max principle we have

$$(13) \quad \lambda_1(M) = \inf \left\{ E(f) \mid f \in C_0^\infty(M), \int_M |f|^2 \omega = 1 \right\}.$$

2.2.1. *A geometric definition.* For any $f \in C^\infty(M)$, define the *horizontal gradient* $\nabla_H f \in \Gamma(\mathcal{D})$ as the horizontal direction of steepest increase of f , that is

$$g(\nabla_H f, X) = df(X), \quad \forall X \in \Gamma(\mathcal{D}).$$

Moreover, the divergence $\operatorname{div}_\omega(X)$ of a smooth vector field $X \in \mathcal{D}$ is

$$\operatorname{div}_\omega(X)\omega = \mathcal{L}_X\omega,$$

where \mathcal{L} denotes the Lie derivative. Thus, Stokes theorem gives

$$\int_M (-\operatorname{div}_\omega(\nabla_H g))f\omega = \int_M g(\nabla_H f, \nabla_H g)\omega, \quad \forall f, g \in C_0^\infty(M).$$

Since $\|\nabla_H f\|^2 = 2H(df)$ for all $f \in C^\infty(M)$, the above formula shows that

$$\Delta_\omega(f) = \operatorname{div}_\omega(\nabla_H f), \quad \forall f \in C_0^\infty(M).$$

By Hörmander theorem [32], Δ_ω is hypoelliptic, and its principal symbol is twice the Hamiltonian function $2H : T^*M \rightarrow \mathbb{R}$.

3. PRELIMINARY CONSTRUCTIONS

We discuss some preliminary constructions concerning integration on vector bundles that we need for the reduction procedure. In this section $\pi : E \rightarrow M$ is a rank k vector bundle on an n dimensional manifold M . For simplicity we assume M to be oriented and E to be oriented (as a vector bundle). If not, the results below remain true replacing volumes with densities. We use coordinates x on $O \subset M$ and $(p, x) \in \mathbb{R}^k \times \mathbb{R}^n$ on $U = \pi^{-1}(O)$ such that the fibers are $E_{q_0} = \{(p, x_0) \mid p \in \mathbb{R}^k\}$. In a compact notation we write, in coordinates, $dp = dp_1 \wedge \dots \wedge dp_k$ and $dx = dx_1 \wedge \dots \wedge dx_n$.

3.1. Vertical volume forms. Consider the fibers $E_q \subset E$ as embedded submanifolds of dimension k . For each $\lambda \in E_q$, let $\Lambda^k(T_\lambda E_q)$ be the space of alternating multi-linear functions on $T_\lambda E_q$. The space

$$\Lambda_v^k(E) := \bigsqcup_{\lambda \in E} \Lambda^k(T_\lambda E_{\pi(\lambda)})$$

defines a rank 1 vector bundle $\Pi : \Lambda_v^k(E) \rightarrow E$, such that $\Pi(\eta) = \lambda$ if $\eta \in \Lambda^k(T_\lambda E_{\pi(\lambda)})$.

To see this, choose coordinates $(p, x) \in \mathbb{R}^k \times \mathbb{R}^n$ on $U = \pi^{-1}(O)$ such that the fibers are $E_{q_0} = \{(p, x_0) \mid p \in \mathbb{R}^k\}$. Thus the vectors $\partial_{p_1}, \dots, \partial_{p_k}$ tangent to the fibers E_q are well defined. The map $\Psi : \Pi^{-1}(U) \rightarrow U \times \mathbb{R}$, defined by $\Psi(\eta) = (\Pi(\eta), \eta(\partial_{p_1}, \dots, \partial_{p_k}))$ is a bijection. Suppose that (U', p', x') is another chart, and similarly $\Psi' : \Pi^{-1}(U') \rightarrow U' \times \mathbb{R}$. Then, on $\Pi^{-1}(U' \cap U) \times \mathbb{R}$ we have $\Psi' \circ \Psi^{-1}(\lambda, \alpha) = (\lambda, \det(\partial q' / \partial q))$. Finally, we apply the vector bundle construction Lemma [36, Lemma 5.5].

Definition 1. A smooth, strictly positive section $\nu \in \Gamma(\Lambda_v^k(E))$ is called a *vertical volume form on E* . In particular, the restriction $\nu_q := \nu|_{E_q}$ of a vertical volume form defines a measure on each fiber E_q .

Lemma 10 (Disintegration 1). *Fix a volume form $\Omega \in \Lambda^{n+k}(E)$ and a volume form $\omega \in \Lambda^n(M)$ on the base space. Then there exists a unique vertical volume form $\nu \in \Lambda_v^k(E)$ such that, for any measurable set $D \subseteq E$ and measurable $f : D \rightarrow \mathbb{R}$,*

$$(14) \quad \int_D f \Omega = \int_{\pi(D)} \left[\int_{D_q} f_q \nu_q \right] \omega(q), \quad f_q := f|_{E_q}, \quad D_q := E_q \cap D.$$

If, in coordinates, $\Omega = \Omega(p, x) dp \wedge dx$ and $\omega = \omega(x) dx$, then

$$\nu|_{(p,x)} = \frac{\Omega(p, x)}{\omega(x)} dp.$$

Proof. The last formula does not depend on the choice of coordinates (p, x) on E . So we can use this as a definition for ν . Moreover, in coordinates,

$$\Omega|_{(p,x)} = \Omega(p, x) dp \wedge dx = \left(\frac{\Omega(p, x)}{\omega(x)} dp \right) \wedge (\omega(x) dx).$$

Both uniqueness and (14) follow from the definition of integration on manifolds and Fubini theorem. \square

3.2. Vertical surface forms. Let $E' \subset E$ be a corank 1 sub-bundle of $\pi : E \rightarrow M$. That is, a submanifold $E' \subset E$ such that $\pi|_{E'} : E' \rightarrow M$ is a bundle, and the fibers $E'_q := \pi^{-1}(q) \cap E' \subset E_q$ are diffeomorphic to a smooth hypersurface $C \subset \mathbb{R}^k$. As a matter of fact, we will only consider the cases in which C is a cylinder or a sphere.

Fix a smooth volume form $\Omega \in \Lambda^{n+k}(E)$. The *Euler vector field* is the generator of homogenous dilations on the fibers $\lambda \mapsto e^\alpha \lambda$, for all $\alpha \in \mathbb{R}$. In coordinates (p, x) on E we have $\epsilon = \sum_{i=1}^n p_i \partial_{p_i}$. If ϵ is transverse to E' we induce a volume form on E' by $\mu := \iota_\epsilon \Omega$.

In this setting, a volume form $\mu \in \Lambda^{n+k-1}(E')$ is called a *surface form*. For any vertical volume form $\nu \in \Lambda_v^k(E)$, we define a measure on the fibers E'_q as $\eta_q = \iota_\epsilon \nu|_{E'_q}$. With an abuse of language, we will refer to such measures as *vertical surface forms*.

Lemma 11 (Disintegration 2). *Fix a surface form $\mu = \iota_\epsilon \Omega \in \Lambda^{n+k-1}(E')$ and a volume form $\omega \in \Lambda^n(M)$ on the base space. For any measurable set $D \subseteq E'$ and measurable $f : D \rightarrow \mathbb{R}$,*

$$\int_D f \mu = \int_{\pi(D)} \left[\int_{D_q} f_q \eta_q \right] \omega(q), \quad f_q := f|_{E'_q}, \quad D_q := E'_q \cap D.$$

Here, $\eta_q = \iota_\epsilon \nu|_{E'_q}$ and ν is the vertical volume form on E defined in Lemma 10.

Proof. Choose coordinates (p, x) on E . As in the proof of Lemma 10

$$\begin{aligned} \mu|_{(p,x)} &= \iota_\epsilon \Omega|_{(p,x)} = \Omega(p, x) \left(\iota_\epsilon dp \wedge dx + (-1)^k dp \wedge \iota_\epsilon dx \right) = \Omega(p, x) \iota_\epsilon dp \wedge dx \\ &= \left(\frac{\Omega(p, x)}{\omega(x)} \iota_\epsilon dp \right) \wedge (\omega(x) dx). \end{aligned}$$

Thus the integral formula above holds with $\eta|_{(p,x)} = \frac{\Omega(p,x)}{\omega(x)} \iota_\epsilon dp$. This, together with the local expression of ν in Lemma 10, gives that $\eta = \iota_\epsilon \nu$. \square

Example 1 (The unit cotangent bundle). We apply the above constructions to $E = T^*M$ and $E' = U^*M$. In this case $E'_q = U_q^*M$ are diffeomorphic to cylinders (or spheres, in the Riemannian case). Moreover, we set $\Omega = \Theta$, the Liouville volume form, and $\mu = \iota_\epsilon \Theta$, the Liouville surface form.⁴ One can check that $\Theta = d\mu$.

Let $\nu \in \Lambda_v^n(T^*M)$ and $\eta = \iota_\epsilon \nu$ as in Lemmas 10 and 11. In canonical coordinates, $\Theta = dp \wedge dx$. Then, if $\omega = \omega(x) dx$,

$$\nu = \frac{1}{\omega(x)} dp \quad \text{and} \quad \eta = \frac{1}{\omega(x)} \sum_{i=1}^n (-1)^{i-1} p_i dp_1 \wedge \dots \wedge \widehat{dp_i} \wedge \dots \wedge dp_n.$$

Choose coordinates x around $q_0 \in M$ such that $\partial_{x_1}|_{q_0}, \dots, \partial_{x_k}|_{q_0}$ is an orthonormal basis for the sub-Riemannian distribution \mathcal{D}_{q_0} . In the associated canonical coordinates we have

$$U_{q_0}^* M = \{(p, x_0) \in \mathbb{R}^{2n} \mid p_1^2 + \dots + p_k^2 = 1\} \simeq \mathbb{S}^{k-1} \times \mathbb{R}^{n-k}.$$

In this chart, η_{q_0} is the $(n-1)$ -volume form of the above infinite cylinder times $1/\omega(x_0)$.

⁴ Let $\vartheta \in \Lambda^1(T^*M)$ be the *tautological form* $\vartheta(\lambda) := \pi^*(\lambda)$. The *Liouville invariant volume* $\Theta \in \Lambda^{2n}(T^*M)$ is $\Theta := (-1)^{\frac{n(n-1)}{2}} d\vartheta \wedge \dots \wedge d\vartheta$. In canonical coordinates (p, x) on T^*M we have $\Theta = dp \wedge dx$.

Remark 4. This construction gives a canonical way to define a measure on U^*M and its fibers in the general sub-Riemannian case, depending only on the choice of the volume ω on the manifold M . It turns out that this measure is also invariant under the Hamiltonian flow. Notice though that in the sub-Riemannian setting, fibers have infinite volume.

3.3. Invariance. Here we focus on the case of interest where $E \subseteq T^*M$ is a rank k vector sub-bundle and $E' \subset E$ is a corank 1 sub-bundle as defined in Section 3.2. We stress that E' is not necessarily a vector sub-bundle, but typically its fibers are cylinders or spheres.

Recall that the sub-Riemannian geodesic flow $\phi_t : T^*M \rightarrow T^*M$ is the Hamiltonian flow of $H : T^*M \rightarrow \mathbb{R}$. Moreover, in our picture, $M \subset N$ is a compact submanifold with boundary ∂M of a larger manifold N , with $\dim M = \dim N = n$.

Definition 2. A sub-bundle $E \subseteq T^*M$ is *invariant* if $\phi_t(\lambda) \in E$ for all $\lambda \in E$ and t such that $\phi_t(\lambda) \in T^*M$ is defined. A volume form $\Omega \in \Lambda^{n+k}(E)$ is *invariant* if $\mathcal{L}_{\vec{H}}\Omega = 0$.

Our definition includes the case of interest for Santaló formula, where sub-Riemannian geodesics may cross $\partial M \neq \emptyset$. In other words, E is invariant if the only way to escape from E through the Hamiltonian flow is by crossing the boundary $\pi^{-1}(\partial M)$. Moreover, if Ω is an invariant volume on an invariant sub-bundle E , then $\phi_t^*\Omega = \Omega$.

Lemma 12 (Invariant induced measures). *Let $E \subseteq T^*M$ be a rank k invariant vector bundle with an invariant volume Ω . Let $E' \subset E$ be a corank 1 invariant sub-bundle. Let \mathfrak{e} be a vector field transverse to E' and $\mu = \iota_{\mathfrak{e}}\Omega$ the induced surface form on E' . Then μ is invariant if and only if $[\vec{H}, \mathfrak{e}]$ is tangent to E' .*

In Example 1, $E = T^*M$ and $E' = U^*M$ are clearly invariant; in particular \vec{H} is tangent to E' . By Liouville theorem, $\Omega = \Theta$ is invariant for any Hamiltonian flow. Moreover, if the Hamiltonian H is homogeneous of degree d (on fibers), one checks that $[\vec{H}, \mathfrak{e}] = -(d-1)\vec{H}$ and Lemma 12 yields the invariance of the Liouville surface measure $\mu = \iota_{\mathfrak{e}}\Theta$. In particular this holds Riemannian and sub-Riemannian geometry, with $d = 2$.

4. SANTALÓ FORMULA

4.1. Assumptions on the boundary. Let (N, \mathcal{D}, g) be a smooth connected sub-Riemannian manifold, of dimension n , without boundary. We focus on a compact n -dimensional submanifold M with piecewise C^2 boundary ∂M .

Let $q \in \partial M$ such that the tangent space is well defined. We say that q is a *characteristic point* if $\mathcal{D}_q \subseteq T_q\partial M$. If q is *non-characteristic*, the *horizontal normal* at q is the unique inward pointing unit vector $\mathbf{n}_q \in \mathcal{D}_q$ orthogonal to $T_q\partial M \cap \mathcal{D}_q$. If $q \in \partial M$ is characteristic, we set $\mathbf{n}_q = 0$. We call $C(\partial M)$ the set of characteristic points. Under the piecewise C^2 regularity of the boundary, we have the following.

Proposition 13. *The set of characteristic points $C(\partial M) \subset \partial M$ is negligible.*

Proof. Without loss of generality we can assume $N = \mathbb{R}^n$ and $\partial M \simeq \{x \in \mathbb{R}^n \mid u(x) = 0\}$ to be the regular level set of some $u \in C^2(\mathbb{R}^n)$. Let us assume by contradiction that $C(\partial M)$ has positive measure. By definition of characteristic point, any $X, Y \in \Gamma(\mathcal{D})$ is tangent to ∂M on $C(\partial M)$. In particular,

$$(15) \quad Xu = Yu = 0 \quad \text{on } C(\partial M).$$

We now claim that (15) implies $XYu = YXu = 0$ a.e. on $C(\partial M)$. Indeed, letting $\nabla(Yu)$ be the Euclidean gradient of $Yu \in C^1(\mathbb{R}^n)$, as a consequence of the Lebesgue density theorem and a first order Taylor expansion we have

$$|\nabla(Yu)|(x) = \lim_{\varepsilon \downarrow 0} \frac{1}{|B_\varepsilon(x)|} \int_{B_\varepsilon(x) \cap C(\partial M)} \frac{|Yu(y) - Yu(x)|}{|y - x|} dy = 0, \quad \text{for a.e. } x \in C(\partial M),$$

Writing $XYu(x) = \nabla(Yu)(x) \cdot X(x)$, where $v \cdot w$ is the Euclidean scalar product of $v, w \in \mathbb{R}^n$, proves the claim.

The above implies that the bracket $[X, Y]$ is tangent to ∂M a.e. on $C(\partial M)$ for any $X, Y \in \Gamma(\mathcal{D})$. In particular, this contradicts the bracket-generating assumption (11). \square

4.2. (Sub-)Riemannian Santaló formula. For any covector $\lambda \in U_q^*M$, the *exit length* $\ell(\lambda)$ is the first time $t \geq 0$ at which the corresponding geodesic $\gamma_\lambda(t) = \pi \circ \phi_t(\lambda)$ leaves M crossing its boundary, while $\tilde{\ell}(\lambda)$ is the smallest between the exit and the cut length along $\gamma_\lambda(t)$. Namely

$$\begin{aligned} \ell(\lambda) &= \sup\{t \geq 0 \mid \gamma_\lambda(t) \in M\}, \\ \tilde{\ell}(\lambda) &= \sup\{t \leq \ell(\lambda) \mid \gamma_\lambda|_{[0,t]} \text{ is minimizing}\}. \end{aligned}$$

We also introduce the following subsets of the unit cotangent bundle $\pi : U^*M \rightarrow M$:

$$\begin{aligned} U^+\partial M &= \{\lambda \in U^*M|_{\partial M} \mid \langle \lambda, \mathbf{n} \rangle > 0\}, \\ U^\star M &= \{\lambda \in U^*M \mid \ell(-\lambda) < +\infty\}, \\ \tilde{U}^\star M &= \{\lambda \in U^\star M \mid \tilde{\ell}(-\lambda) = \ell(-\lambda)\}. \end{aligned}$$

Some comments are in order. The set $U^+\partial M$ consists of the unit covectors $\lambda \in \pi^{-1}(\partial M)$ such that the associated geodesic enters the set M for arbitrary small $t > 0$. The *visible set* $U^\star M$ is the set of covectors that can be reached (in finite time) starting from $\pi^{-1}(\partial M)$ and following the geodesic flow. If we restrict to covectors that can be reached *optimally* in finite time, we obtain the *optimally visible set* $\tilde{U}^\star M$ (see Fig. 1).

Lemma 14. *The cut-length $c : U^*M \rightarrow (0, +\infty]$ is upper semicontinuous (and hence measurable). Moreover, if any couple of points in M can be joined by a minimizing non-abnormal geodesic, c is continuous.*

Proof. The result follows as in [19, Theorem III.2.1]. We stress that the key part of the proof of the second statement is the fact that, in absence of abnormal minimizers, a point is in the cut locus of another if and only if (i) it is conjugate along some minimizing geodesic or (ii) there exist two distinct minimizing geodesics joining them. \square

Lemma 15. *The map $\ell : U^+\partial M \rightarrow (0, +\infty]$ is lower semicontinuous (and hence measurable). Moreover, $\tilde{\ell} : U^+\partial M \rightarrow (0, +\infty]$ is measurable.*

Proof. Let $\lambda_0 \in U^+\partial M$. Consider a sequence λ_n such that $\liminf_{\lambda \rightarrow \lambda_0} \ell(\lambda) = \lim_n \ell(\lambda_n)$. Then, the trajectories $\gamma_n(t) = \pi \circ \phi_t(\lambda_n)$ for $t \in [0, \ell(\lambda_n)]$ converge uniformly as $n \rightarrow +\infty$ to the trajectory $\gamma_0(t) = \phi_t(\lambda_0)$ for $t \in [0, \delta]$ where $\delta = \lim_n \ell(\lambda_n)$. Moreover, by continuity of ∂M and the fact that $\gamma_n(\ell(\lambda_n)) \in \partial M$, it follows that $\gamma_0(\delta) \in \partial M$. This proves that $\delta \geq \ell(\lambda_0)$, proving the first part of the statement.

To complete the proof, observe that $\tilde{\ell} = \min\{\ell, c\}$, which are measurable by the previous claim and Lemma 14. \square

Fix a volume form ω on M (or density, if M is not orientable). In any case, ω and $\sigma := \iota_{\mathbf{n}}\omega$ induce positive measures on M and ∂M , respectively. According to Lemmas 10 and 11, these induce measures ν_q and $\eta_q = \iota_{\mathbf{e}_q}\nu_q$ on T_q^*M and U_q^*M , respectively.

Theorem 16 (Santaló formulas). *The visible set $U^\star M$ and the optimally visible set $\tilde{U}^\star M$ are measurable. Moreover, for any measurable function $F : U^*M \rightarrow \mathbb{R}$ we have*

$$(16) \quad \begin{aligned} \int_{U^\star M} F \mu &= \int_{\partial M} \left[\int_{U_q^+\partial M} \left(\int_0^{\ell(\lambda)} F(\phi_t(\lambda)) dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q(\lambda) \right] \sigma(q), \\ \int_{\tilde{U}^\star M} F \mu &= \int_{\partial M} \left[\int_{U_q^+\partial M} \left(\int_0^{\tilde{\ell}(\lambda)} F(\phi_t(\lambda)) dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q(\lambda) \right] \sigma(q). \end{aligned}$$

Remark 5. Even if M is compact and hence $\tilde{\ell} < +\infty$, in general $\tilde{U}^{\check{\ast}}M \subsetneq U^{\check{\ast}}M$. Moreover, if also $\ell < +\infty$ (that is, all geodesics reach the boundary of M in finite time), then $U^{\check{\ast}}M = U^*M$. Thus, our statement of Santaló formula contains [19, Theorem VII.4.1].

Remark 6. If ∂M is only Lipschitz and $C(\partial M)$ has positive measure, the above Santaló formulas still hold by removing on the left hand side from $U^{\check{\ast}}M$ and $\tilde{U}^{\check{\ast}}M$ the set $\{\phi_t(\lambda) \mid \pi(\lambda) \in C(\partial M) \text{ and } t \geq 0\}$. Nothing changes on the right hand side as $\sigma(C(\partial M)) = 0$ by definition of \mathbf{n} .

Proof. Let $A \subset [0, +\infty) \times U^+\partial M$ be the set of pairs (t, λ) such that $0 < t < \ell(\lambda)$. By Lemma 15 it follows that A is measurable. Let also $Z = \pi^{-1}(\partial M) \subset U^{\check{\ast}}M$ which clearly has zero measure in U^*M . Define $\phi : A \rightarrow U^{\check{\ast}}M \setminus Z$ as $\phi(t, \lambda) = \phi_t(\lambda)$. This is a smooth diffeomorphism, whose inverse is $\phi^{-1}(\bar{\lambda}) = (-\phi_{\ell(-\bar{\lambda})}(-\bar{\lambda}), \ell(-\bar{\lambda}))$. In particular, $U^{\check{\ast}}M$ is measurable. Then, using Lemma 17 (see below),

$$(17) \quad \int_{U^{\check{\ast}}M} F \mu = \int_{\phi(A)} F \mu = \int_A (F \circ \phi) \phi^* \mu = \\ = \int_{\partial M} \left[\int_{U_q^+\partial M} \left(\int_0^{\ell(\lambda)} F(\phi_t(\lambda)) dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q(\lambda) \right] \sigma(q).$$

by Fubini Theorem. Analogously, with $\tilde{A} = \{(t, \lambda) \mid 0 < t < \tilde{\ell}(\lambda)\}$ and $\tilde{Z} = Z \cup \{\phi_{\tilde{\ell}(\lambda)}(\lambda) \mid \lambda \in U^+\partial M\}$ the map $\phi : \tilde{A} \rightarrow \tilde{U}^{\check{\ast}}M \setminus \tilde{Z}$ is a diffeomorphism with the same inverse. Then, the same computations as (17) replacing A with \tilde{A} and Z with \tilde{Z} yield (16). \square

Lemma 17. *The following local identity of elements of $\Lambda^{2n-1}(\mathbb{R} \times U^+\partial M)$ holds*

$$\phi^* \mu|_{(t,\lambda)} = \langle \lambda, \mathbf{n}_q \rangle dt \wedge \sigma \wedge \eta, \quad \lambda \in U^+\partial M,$$

where, in canonical coordinates (p, x) on T^*M

$$\eta = \iota_{\mathbf{e}} \nu, \quad \nu = \frac{1}{\omega(x)} dp, \quad \sigma = \iota_{\mathbf{n}} \omega, \quad \omega = \omega(x) dx.$$

Proof. For any $(t, \lambda) \in \mathbb{R} \times U^+\partial M$ let $\{\partial_t, v_1, \dots, v_{2n-2}\}$ be a set of independent vectors in $T(\mathbb{R} \times U^+\partial M) = T\mathbb{R} \oplus TU^+\partial M$. Observe that $\phi^* \mu = dt \wedge (\iota_{\partial_t} \phi^* \mu)$. Then,

$$\iota_{\partial_t} \phi^* \mu(v_1, \dots, v_{2n-2}) = \mu|_{\phi(t,\lambda)} \left(d_{(t,\lambda)} \phi \partial_t, d_{(t,\lambda)} \phi v_1, \dots, d_{(t,\lambda)} \phi v_{2n-2} \right).$$

Notice that,

- (a) $d_{(t,\lambda)} \phi v_i = (d_\lambda \phi_t) v_i$ for any $i = 1, \dots, 2n-2$,
- (b) $d_{(t,\lambda)} \phi \partial_t = (d_\lambda \phi_t) \vec{H}$, this is in fact just $\vec{H}|_{\phi_t(\lambda)}$.

Now it follows that

$$(18) \quad \iota_{\partial_t} \phi^* \mu = \iota_{\vec{H}} \phi_t^* \mu = \iota_{\vec{H}} \mu,$$

where in the last passage we used the invariance of μ (see the discussion below Lemma 12). By Lemma 11 and its proof (in particular see Example 1) locally $\mu = \eta \wedge \omega$. By the properties of the interior product,

$$(19) \quad \iota_{\vec{H}} \mu = (\iota_{\vec{H}} \eta) \wedge \omega + (\iota_{\vec{H}} \omega) \wedge \eta.$$

The first term on the r.h.s. vanishes: as a $2n-2$ form, its value at a point $\lambda \in U^+\partial M$ is completely determined by its action on $2n-2$ independent vectors of $T_\lambda U^+\partial M$. We can choose coordinates such that $\partial M = \{x_n = 0\}$. Then a basis of $T_\lambda U^+\partial M$ is given by $\partial_{x_1}, \dots, \partial_{x_{n-1}}$ and a set of $n-1$ vectors $v_i = \sum_{j=1}^n v_i^j \partial_{p_j}$ in $T_\lambda U_{\pi(\lambda)}^+ \partial M$. Since $\iota_{\vec{H}} \eta$ is a $n-2$ form, then ω necessarily acts on at least one v_i , and vanishes. Now, notice that

$$(20) \quad \iota_{\vec{H}} \omega|_\lambda(\cdot) = \omega|_{\pi(\lambda)}(\pi_* \vec{H}, \pi_* \cdot) = \langle \lambda, \mathbf{n}_{\pi(\lambda)} \rangle \omega|_{\pi(\lambda)}(\mathbf{n}_{\pi(\lambda)}, \pi_* \cdot) = \langle \lambda, \mathbf{n}_{\pi(\lambda)} \rangle \sigma|_\lambda(\cdot).$$

Putting together (18), (19), and (20) completes the proof of the statement. \square

4.3. Reduced Santaló formula. The following reduction procedure replaces the non-compact set $U^{\check{c}}M$ in Theorem 16 with a compact subset that we now describe.

To carry out this procedure we fix a transverse sub-bundle $\mathcal{V} \subset TM$ such that $TM = \mathcal{D} \oplus \mathcal{V}$. We assume that \mathcal{V} is the orthogonal complement of \mathcal{D} w.r.t. to a Riemannian metric g such that $g|_{\mathcal{D}}$ coincides with the sub-Riemannian one and the associated Riemannian volume coincides with ω . In the Riemannian case, where \mathcal{V} is trivial, this forces $\omega = \omega_R$, the Riemannian volume. In the genuinely sub-Riemannian case there is no loss of generality since this assumption is satisfied for any choice of ω .

Definition 3. The *reduced cotangent bundle* is the rank k vector bundle $\pi : T^*M^r \rightarrow M$ of covectors that annihilate the vertical directions:

$$T^*M^r := \{\lambda \in T^*M \mid \langle \lambda, v \rangle = 0 \text{ for all } v \in \mathcal{V}\}.$$

The *reduced unit cotangent bundle* is $U^*M^r := U^*M \cap T^*M^r$.

Observe that U^*M^r is a corank 1 sub-bundle of T^*M^r , whose fibers are spheres \mathbb{S}^{k-1} . If T^*M^r is invariant in the sense of Definition 2, we can apply the construction of Section 3.3. The Liouville volume Θ on T^*M induces a volume on T^*M^r as follows.

Let X_1, \dots, X_k and Z_1, \dots, Z_{n-k} be local orthonormal frames for \mathcal{D} and \mathcal{V} , respectively. Let $u_i(\lambda) := \langle \lambda, X_i \rangle$ and $v_j(\lambda) := \langle \lambda, Z_j \rangle$ smooth functions on T^*M . Thus

$$T^*M^r = \{\lambda \in T^*M \mid v_1(\lambda) = \dots = v_{n-k}(\lambda) = 0\}.$$

For all $q \in M$ where the fields are defined, $(u, v) : T_q^*M \rightarrow \mathbb{R}^n$ are smooth coordinates on the fiber and hence $\partial_{u_1}, \dots, \partial_{u_k}, \partial_{v_1}, \dots, \partial_{v_{n-k}}$ are vectors on $T_\lambda(T_q^*M) \subset T_\lambda(T^*M)$ for all $\lambda \in \pi^{-1}(q)$. In particular, the vector fields $\partial_{v_1}, \dots, \partial_{v_{n-k}}$ are transverse to T^*M^r , hence we give the following definition.

Definition 4. The *reduced Liouville volume* $\Theta^r \in \Lambda^{n+k}(T^*M^r)$ is

$$\Theta_\lambda^r := \Theta_\lambda(\underbrace{\dots, \dots, \dots}_{k \text{ vectors}}, \partial_{v_1}, \dots, \partial_{v_{n-k}}, \underbrace{\dots, \dots, \dots}_{n \text{ vectors}}), \quad \forall \lambda \in T^*M^r.$$

The above definition of Θ^r does not depend on the choice of the local orthonormal frame $\{X_1, \dots, X_k, Z_1, \dots, Z_{n-k}\}$ and Riemannian metric $g|_{\mathcal{V}}$ on the complement, as long as its Riemannian volume remains the fixed one, ω . In fact, let X', Z' be a different frame for a different Riemannian metric $g'|_{\mathcal{V}}$. Then⁵, $X' = RX$ and $Z' = SX + TZ$ for $R \in \text{SO}(k)$, $T \in \text{SL}(n-k)$ and $S \in \text{M}(k, n)$. One can check that $\partial_v = S\partial_{v'} + T\partial_{v''}$ and that $\Omega^r = \Omega(\dots, \partial_v, \dots) = \Omega(\dots, T\partial_{v'}, \dots) = (\Omega^r)'$, where both frames are defined.

Assumptions for the reduction. We assume the following hypotheses:

- (H1) The bundle $T^*M^r \subseteq T^*M$ is invariant.
- (H2) The *reduced Liouville volume* is invariant, i.e. $\mathcal{L}_{\vec{H}}\Theta^r = 0$.

Remark 7. Assumption (H1) depends only on \mathcal{V} , while (H2) depends also on ω (since Θ^r does). In particular, in the Riemannian case, both are trivially satisfied.

Under these assumptions $U^*M^r = U^*M \cap T^*M^r$ is an invariant corank 1 sub-bundle of T^*M^r . Moreover, $\mu^r = \iota_{\mathbf{e}}\Theta^r$ is an invariant surface form on U^*M^r . This follows from Lemma 12 observing that $[\vec{H}, \mathbf{e}] = -\vec{H}$ is tangent to U^*M^r . As in Section 3.1, the volume $\Theta^r \in \Lambda^{n+k}(T^*M^r)$ induces a vertical volume ν_q^r on the fibers $T_q^*M^r$ and a vertical surface form $\eta_q^r = \iota_{\mathbf{e}}\nu_q^r$ on $U_q^*M^r$. As a consequence of Lemmas 10 and 11 the latter has the following explicit expression, whose proof is straightforward.

⁵For simplicity, assume that \mathcal{D} is orientable as a vector bundle and that X_1, \dots, X_k is an oriented frame.

Lemma 18 (Explicit reduced vertical measure). *Let $q_0 \in M$ and fix a set of canonical coordinates (p, x) such that q_0 has coordinates x_0 and*

- $\{\partial_{x_1}, \dots, \partial_{x_k}\}_{q_0}$ is an orthonormal basis of \mathcal{D}_{q_0} ,
- $\{\partial_{x_{k+1}}, \dots, \partial_{x_n}\}_{q_0}$ is an orthonormal basis of \mathcal{V}_{q_0} .

In these coordinates $\omega|_{x_0} = dx|_{q_0}$. Then $\nu_{q_0}^r = \text{vol}_{\mathbb{R}^k}$ and $\eta_{q_0}^r = \text{vol}_{\mathbb{S}^{k-1}}$. In particular,

$$\int_{U_{q_0}^* M^r} \eta_{q_0}^r = |\mathbb{S}^{k-1}|, \quad \forall q_0 \in M,$$

where $|\mathbb{S}^{k-1}|$ denotes the Lebesgue measure of \mathbb{S}^{k-1} and $\text{vol}_{\mathbb{R}^k}, \text{vol}_{\mathbb{S}^{k-1}}$ denote the Euclidean volume forms of \mathbb{R}^k and \mathbb{S}^{k-1} .

We now state the reduced Santaló formulas. The sets $U^+ \partial M^r$, $U^\star M^r$, and $\tilde{U}^\star M^r$ are defined from their unreduced counterparts by taking the intersection with $T^* M^r$.

Theorem 19 (Reduced Santaló formulas). *The visible set $U^\star M^r$ and the optimally visible set $\tilde{U}^\star M^r$ are measurable. For any measurable function $F : U^* M^r \rightarrow \mathbb{R}$ we have*

$$(21) \quad \begin{aligned} \int_{U^\star M^r} F \mu^r &= \int_{\partial M} \left[\int_{U_q^+ \partial M^r} \left(\int_0^{\ell(\lambda)} F(\phi_t(\lambda)) dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q^r(\lambda) \right] \sigma(q), \\ \int_{\tilde{U}^\star M^r} F \mu^r &= \int_{\partial M} \left[\int_{U_q^+ \partial M^r} \left(\int_0^{\tilde{\ell}(\lambda)} F(\phi_t(\lambda)) dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q^r(\lambda) \right] \sigma(q). \end{aligned}$$

Proof. The proof follows the same steps as the one of Theorem 16 replacing the invariant sub-bundles, volumes, and surface forms with their reduced counterparts. \square

Remark 8. Let H_{sR} be the sub-Riemannian Hamiltonian and H_R be the Riemannian Hamiltonian of the Riemannian extension. The two Hamiltonians are (locally on $T^* M$)

$$H_R = \frac{1}{2} \left(\sum_{i=1}^k u_i^2 + \sum_{j=1}^{n-k} v_j^2 \right), \quad H_{sR} = \frac{1}{2} \sum_{i=1}^k u_i^2.$$

Let $\phi_t^{sR} = e^{t\tilde{H}_{sR}}$ and $\phi_t^R = e^{t\tilde{H}_R}$ be their Hamiltonian flows. Since $T^* M^r = \{\lambda \mid v_1(\lambda) = \dots = v_{n-k}(\lambda) = 0\}$, by assumption **(H1)** we have

$$H_{sR} = H_R, \quad \text{and} \quad \phi_t^{sR} = \phi_t^R \quad \text{on } T^* M^r.$$

In particular, the sub-Riemannian geodesics with initial covector $\lambda \in U^* M^r$ are also geodesics of the Riemannian extension and viceversa.

5. EXAMPLES

5.1. Carnot groups. A *Carnot group* (G, \star) of step m is a connected, simply connected Lie group of dimension n , such that its Lie algebra $\mathfrak{g} = T_e G$ admits a nilpotent stratification of step m , that is

$$\mathfrak{g} = \mathfrak{g}_1 \oplus \dots \oplus \mathfrak{g}_m,$$

with

$$[\mathfrak{g}_1, \mathfrak{g}_j] = \mathfrak{g}_{i+j}, \quad \forall 1 \leq j \leq m, \quad \mathfrak{g}_m \neq \{0\}, \quad \mathfrak{g}_{m+1} = \{0\}.$$

Let \mathcal{D} be the left-invariant distribution generated by \mathfrak{g}_1 , and consider any left-invariant sub-Riemannian structure on G induced by a scalar product on \mathfrak{g}_1 .

We identify $G \simeq \mathbb{R}^n$ with a polynomial product law by choosing a basis for \mathfrak{g} as follows. Recall that the group exponential map,

$$\exp_G : \mathfrak{g} \rightarrow G,$$

associates with $V \in \mathfrak{g}$ the element $\gamma(1)$, where $\gamma : [0, 1] \rightarrow G$ is the unique integral line starting from $\gamma(0) = 0$ of the left invariant vector field associated with V . Since G is simply connected and \mathfrak{g} is nilpotent, \exp_G is a smooth diffeomorphism.

Let $d_j := \dim \mathfrak{g}_j$. Indeed $d_1 = k$. Let $\{X_i^j\}$, for $j = 1, \dots, m$ and $i = 1, \dots, d_j$ be an adapted basis, that is $\mathfrak{g}_j = \text{span}\{X_1^j, \dots, X_{d_j}^j\}$. In exponential coordinates we identify

$$(x^1, \dots, x^m) \simeq \exp_G \left(\sum_{j=1}^m \sum_{i=1}^{d_j} x_i^j X_i^j \right), \quad x^j \in \mathbb{R}^{d_j}.$$

The identity $e \in G$ is the point $(0, \dots, 0) \in \mathbb{R}^n$ and, by the Baker-Campbell-Hausdorff formula the group law \star is a polynomial expression in the coordinates (x^1, \dots, x^m) . Finally,

$$X_i^j = \left. \frac{\partial}{\partial x_i^j} \right|_0,$$

so that $\mathcal{D}|_e \simeq \{(x, 0, \dots, 0) \mid x \in \mathbb{R}^k\}$ and $\mathcal{D}_q = L_{q\star} \mathcal{D}|_e$, where $L_{q\star}$ is the differential of the left-translation $L_q(p) := q \star p$.

We equip G with its Haar measure that is proportional to the Lebesgue volume of \mathbb{R}^n . In order to apply the reduction procedure of Section 4.3, let \mathcal{V} be the left-invariant distribution generated by

$$\mathcal{V}|_e := \mathfrak{g}_2 \oplus \dots \oplus \mathfrak{g}_m,$$

and consider any left-invariant scalar product $g|_{\mathcal{V}}$ on \mathcal{V} . Thus, up to a renormalization, $g = g|_{\mathcal{D}} \oplus g|_{\mathcal{V}}$ is a left-invariant Riemannian extension such that $TM = \mathcal{D} \oplus \mathcal{V}$ is an orthogonal direct sum and its Riemannian volume coincides with the Lebesgue one.

Proposition 20. *Any Carnot group satisfies assumptions (H1) and (H2).*

Proof. Let $X_1, \dots, X_k \in \Gamma(\mathcal{D})$ and $Z_1, \dots, Z_{n-k} \in \Gamma(\mathcal{V})$ be a global frame of left-invariant orthonormal vector fields. Let $u_i(\lambda) := \langle \lambda, X_i \rangle$ and $v_j(\lambda) := \langle \lambda, Z_j \rangle$ be smooth functions on T^*G . We have the following expressions for the Poisson brackets

$$\{u_i, v_j\} = \sum_{i=1}^k \sum_{\ell=1}^{n-k} d_{ij}^{\ell} v_{\ell}, \quad i = 1, \dots, k, \quad j = 1, \dots, n-k,$$

for some constants d_{ij}^{ℓ} . We stress that the above expression does not depend on the u_i 's, as a consequence of the graded structure. Denoting the derivative along the integral curves of \vec{H} with a dot, we have

$$\dot{v}_j = \{H, v_j\} = \sum_{i=1}^k u_i \{u_i, v_j\} = \sum_{i=1}^k \sum_{\ell=1}^{n-k} u_i d_{ij}^{\ell} v_{\ell}.$$

Thus, any integral line of \vec{H} starting from $\lambda \in T^*M^r = \{v_1 = \dots = v_{n-k} = 0\}$ remains in T^*M^r and the latter is invariant.

To prove the invariance of Θ^r , consider, for any fixed left-invariant $X \in \Gamma(\mathcal{D})$, the adjoint map $\text{ad}_X : \mathcal{V}|_e \rightarrow \mathcal{V}|_e$, given by $\text{ad}_X(Z) = [X, Z]|_e$. This map is well defined (as a consequence of the graded structure) and nilpotent. In particular $\text{Trace}(\text{ad}_X) = 0$. Thus, we obtain from an explicit computation (see Appendix A)

$$\mathcal{L}_{\vec{H}} \Theta^r = - \left(\sum_{i=1}^k \sum_{j=1}^{n-k} u_i d_{ij}^j \right) \Theta^r = - \left(\sum_{i=1}^k u_i \text{Trace}(\text{ad}_{X_i}) \right) \Theta^r = 0. \quad \square$$

Proposition 21 (Characterization of reduced geodesics for Carnot groups). *The geodesics $\gamma_{\lambda}(t)$ with initial covector $\lambda \in T_q^*M^r$ are obtained by left-translation of straight lines, that is, in exponential coordinates,*

$$\gamma_{\lambda}(t) = q \star (ut, 0, \dots, 0), \quad u \in \mathbb{R}^k.$$

Proof. Let $X_1, \dots, X_k \in \Gamma(\mathcal{D})$ and $Z_1, \dots, Z_{n-k} \in \Gamma(\mathcal{V})$ be a global frame of left-invariant orthonormal vector fields. Let $u_i(\lambda) := \langle \lambda, X_i \rangle$ and $v_j(\lambda) := \langle \lambda, Z_j \rangle$ be smooth functions on T^*G . The extremal $\lambda(t) = \phi_t(\lambda)$, with initial covector $\lambda = (q, u, 0)$ satisfies $v \equiv 0$ by Proposition 20 and, as a consequence of the graded structure,

$$\dot{u}_i = \{H, u_i\} = \sum_{j=1}^k u_j \{u_j, u_i\} = \sum_{j=1}^k \sum_{\ell=1}^{n-k} u_j c_{ji}^\ell v_\ell = 0,$$

In particular $\lambda(t) = (q(t), u, 0)$ for a constant $u \in \mathbb{R}^k$. Moreover the geodesic $\gamma_\lambda(t) = \pi(\lambda(t))$ satisfies

$$\dot{\gamma}_\lambda(t) = \sum_{i=1}^k u_i X_i(\gamma_\lambda(t)).$$

Since the u_i 's are constants, $\gamma_\lambda(t)$ is an integral curve of $\sum_{i=1}^k u_i X_i$ starting from q . Then $L_q^{-1} \gamma_\lambda(t)$ is an integral curve of $\sum_{i=1}^k u_i L_{q^*}^{-1} X_i = \sum_{i=1}^k u_i X_i$ starting from the identity. By definition of exponential coordinates

$$\gamma_\lambda(t) = q \star \exp_G \left(t \sum_{i=1}^k u_i X_i \right) \simeq q \star (ut, 0). \quad \square$$

Remark 9. In the case of a step 2 Carnot group, the group law is linear when written in exponential coordinates. In fact, for a fixed left-invariant basis $X_1, \dots, X_k \in \Gamma(\mathcal{D})$ and $Z_1, \dots, Z_{n-k} \in \Gamma(\mathcal{V})$ it holds

$$[X_i, X_j] = \sum_{\ell=1}^{n-k} c_{ij}^\ell Z_\ell, \quad c_{ij}^\ell \in \mathbb{R}.$$

By the Baker-Campbell-Hausdorff formula, $(x, z) \star (x', z') = (x' + x, z' + z + f(x, x'))$, where

$$f(x, x')_\ell = \frac{1}{2} \sum_{i,j=1}^k x_i c_{ij}^\ell x'_j, \quad \ell = 1, \dots, n-k.$$

As a consequence, the geodesics $\gamma_\lambda(t)$ with initial covector $\lambda \in T_q^* M^r$ span the set $q \star \mathcal{D}|_e$. The latter is not an hyperplane, in general, when $q \neq e$ and the step $m > 2$.

Example 2 (Heisenberg group). The $(2d+1)$ -dimensional Heisenberg group \mathbb{H}_{2d+1} is the sub-Riemannian structure on \mathbb{R}^{2d+1} where (\mathcal{D}, g) is given by the following set of global orthonormal fields

$$X_i := \partial_{x_i} - \frac{1}{2} \sum_{j=1}^{2d} J_{ij} x_j \partial_z, \quad J = \begin{pmatrix} 0 & \mathbb{I}_n \\ -\mathbb{I}_n & 0 \end{pmatrix}, \quad i = 1, \dots, 2d,$$

written in coordinates $(x, z) \in \mathbb{R}^{2d} \times \mathbb{R}$. The distribution is bracket-generating, as $[X_i, X_j] = J_{ij} \partial_z$. These fields generate a stratified Lie algebra, nilpotent of step 2, with

$$\mathfrak{g}_1 = \text{span}\{X_1, \dots, X_{2d}\}, \quad \mathfrak{g}_2 = \text{span}\{\partial_z\}.$$

There is a unique connected, simply connected Lie group G such that $\mathfrak{g} = \mathfrak{g}_1 \oplus \mathfrak{g}_2$ is its Lie algebra of left-invariant vector fields. The group exponential map $\exp_G : \mathfrak{g} \rightarrow G$ is a smooth diffeomorphism and then we identify $G = \mathbb{R}^{2d+1}$ with the polynomial product law

$$(x, z) \star (x', z') = \left(x + x', z + z' + \frac{1}{2} x^* J x' \right).$$

Notice that X_1, \dots, X_{2d} (and ∂_z) are left-invariant.

To carry on the reduction, we consider the Riemannian extension g such that ∂_z is a unit vector orthogonal to \mathcal{D} . The geodesics associated with $\lambda \in U^* M^r$ and starting from

q reach the whole Euclidean plane $q \star \{z = 0\}$ (the left-translation of $\mathbb{R}^{2d} \subset \mathbb{R}^{2d+1}$). At $q = (x, z)$ this is the plane orthogonal to the vector $(\frac{1}{2}Jx, 1)$ w.r.t. the Euclidean metric.

5.2. Riemannian foliations with bundle like metric. Roughly speaking, a Riemannian foliation has bundle like metric if locally is a Riemannian submersion w.r.t. the projection along the leaves.

Definition 5. Let M be a smooth and connected n -dimensional Riemannian manifold. A codimension k foliation \mathcal{F} on M is said to be *Riemannian with bundle like metric* if there exists a maximal collection of pairs $\{(U_\alpha, \pi_\alpha), \alpha \in I\}$ of open subsets U_α of M and submersions $\pi_\alpha : U_\alpha \rightarrow U_\alpha^0 \subset \mathbb{R}^k$ such that

- $\{U_\alpha\}_{\alpha \in I}$ is a covering of M
- If $U_\alpha \cap U_\beta \neq \emptyset$, there exists a local diffeomorphism $\Psi_{\alpha\beta} : \mathbb{R}^k \rightarrow \mathbb{R}^k$ such that $\pi_\alpha = \Psi_{\alpha\beta} \pi_\beta$ on $U_\alpha \cap U_\beta$
- the maps $\pi_\alpha : U_\alpha \rightarrow U_\alpha^0$ are Riemannian submersions when U_α^0 are endowed with a given Riemannian metric

Moreover the foliation is *totally geodesic* if its leaves are totally geodesic submanifolds [10].

We say that a vector field $X \in \Gamma(TM)$ is basic if, locally on any U_α , it is π_α -related with some vector X^0 on U_α^0 . If $X \in \Gamma(TM)$ is basic, and $V \in \Gamma(\mathcal{V})$ is vertical, then the Lie bracket $[X, V]$ is vertical. In this setting we consider a local orthonormal frame $Z_1, \dots, Z_{n-k} \in \Gamma(\mathcal{V})$ of vertical vector fields and a local orthonormal frame of basic vector fields $X_1, \dots, X_k \in \Gamma(\mathcal{D})$ for the distribution. The structural functions are defined as

$$[X_i, X_j] = \sum_{\ell=1}^k b_{ij}^\ell X_\ell + \sum_{\ell=1}^{n-k} c_{ij}^\ell Z_\ell, \quad [X_i, Z_j] = \sum_{\ell=1}^{n-k} d_{ij}^\ell Z_\ell, \quad [Z_i, Z_j] = \sum_{\ell=1}^{n-k} e_{ij}^\ell Z_\ell.$$

The totally geodesic assumption is equivalent to the fact that any basic horizontal vector field X generates a vertical isometry, that is

$$(22) \quad (\mathcal{L}_X g)(Z, W) = 0, \quad \forall Z, W \in \Gamma(\mathcal{V}) \quad \iff \quad d_{ij}^\ell = -d_{i\ell}^j.$$

To any Riemannian foliation with bundle-like metric we associate the splitting $TM = \mathcal{D} \oplus \mathcal{V}$, where \mathcal{V} is the bundle of vectors tangent to the leaves of the foliation and \mathcal{D} is its orthogonal complement (we call \mathcal{V} the bundle of *vertical directions*, and its sections *vertical vector fields*). If \mathcal{D} is bracket-generating, then $(\mathcal{D}, g|_{\mathcal{D}})$ is indeed a sub-Riemannian structure on M that we refer to as *tamed by a foliation* and we assume to be equipped with the corresponding Riemannian volume.

Proposition 22. *Any sub-Riemannian structure tamed by a foliation satisfies assumptions (H1) and (H2).*

Proof. Locally, T^*M^r is the zero-locus of the functions $v_i(\lambda) = \langle \lambda, Z_i \rangle$ for some family $\{Z_j\}_{j=1}^{n-k}$ of generators of \mathcal{V} . Thus, denoting the derivative along the integral curves of \vec{H} with a dot, we have

$$\dot{v}_j = \{H, v_i\} = \sum_{i=1}^k u_i \{u_i, v_j\} = \sum_{i=1}^k \sum_{\ell=1}^{n-k} u_i d_{ij}^\ell v_\ell = 0, \quad \text{on } T^*M^r.$$

This readily implies the invariance of T^*M^r . To prove the invariance of Θ^r , we obtain from an explicit computation (see Appendix A)

$$\mathcal{L}_{\vec{H}}(\Theta^r) = - \left(\sum_{i=1}^k \sum_{\ell=1}^{n-k} u_i d_{i\ell}^\ell \right) \Theta^r = 0,$$

where, in the last step, we used the totally geodesic assumption (22). \square

5.2.1. *Riemannian submersions.* A Riemannian submersion $\pi : (M, g) \rightarrow (\bar{M}, \bar{g})$ is trivially a Riemannian foliation with bundle-like metric. Let M be a sub-Riemannian manifold tamed by a Riemannian submersion $\pi : M \rightarrow \bar{M}$. We have the following characterization.

Proposition 23. *Let M be a sub-Riemannian manifold tamed by a Riemannian submersion $\pi : M \rightarrow \bar{M}$. Then $\gamma_\lambda : [0, T] \rightarrow M$ is a sub-Riemannian geodesic associated with $\lambda \in U^*M^r$ if and only if it is the lift of a Riemannian geodesic $\bar{\gamma}_\lambda := \pi \circ \gamma_\lambda$ of \bar{M} .*

Proof. Let $\bar{X}_1, \dots, \bar{X}_k \in \Gamma(T\bar{M})$ be a local orthonormal frame for (\bar{M}, \bar{g}) . Let $X_1, \dots, X_k \in \Gamma(\mathcal{D})$ the corresponding local orthonormal frame of basic vector fields on M , such that $\pi_*X_i = \bar{X}_i$. Let $Z_1, \dots, Z_{n-k} \in \Gamma(\mathcal{V})$ be a local orthonormal frame for \mathcal{V} . Indeed

$$[X_i, X_j] = \sum_{\ell=1}^k b_{ij}^\ell X_\ell + \sum_{\ell=1}^{n-k} c_{ij}^\ell Z_\ell, \quad b_{ij}^\ell, c_{ij}^\ell \in C^\infty(M).$$

Since the X_i 's are basic, the functions $b_{ij}^\ell \in C^\infty(M)$ are constant along the fibers of the submersion and descend to well defined functions in $C^\infty(\bar{M})$. Moreover

$$[\bar{X}_i, \bar{X}_j] = \sum_{\ell=1}^k b_{ij}^\ell \bar{X}_\ell.$$

By Proposition 22, sub-Riemannian extremals $\lambda(t) \in U^*M^r$ satisfy

$$(23) \quad \dot{v}_j(t) \equiv 0, \quad \dot{u}_j(t) = \sum_{i,\ell=1}^k u_i(t) b_{ij}^\ell u_\ell(t), \quad \dot{\gamma}_\lambda(t) = \sum_{i=1}^k u_i(t) X_i(\gamma_\lambda(t)),$$

where the structural functions $b_{ij}^\ell = b_{ij}^\ell(\gamma_\lambda(t))$ are computed along the sub-Riemannian geodesic. On the other hand, Riemannian extremals $\bar{\lambda}(t) \in U\bar{M}$ satisfy

$$(24) \quad \dot{\bar{u}}_j(t) = \sum_{i,\ell=1}^k \bar{u}_i(t) b_{ij}^\ell \bar{u}_\ell(t), \quad \dot{\gamma}_\lambda(t) = \sum_{i=1}^k \bar{u}_i(t) \bar{X}_i(\bar{\gamma}_\lambda(t)),$$

where $\bar{u}_i : T^*\bar{M} \rightarrow \mathbb{R}$ are the smooth functions $u_i(\bar{\lambda}) = \langle \bar{\lambda}, \bar{X}_i \rangle$, for $i = 1, \dots, k$ and are computed along the extremal. The statement follows by observing that the projections $\bar{\gamma}_\lambda = \pi \circ \gamma_\lambda$ of sub-Riemannian extremals satisfy (24) with $\bar{u}_i(t) = u_i(t)$. Viceversa, for any Riemannian geodesic $\bar{\gamma}_\lambda$ on \bar{M} , its horizontal lift γ_λ on M satisfies (23) with $u_i(t) = \bar{u}_i(t)$ and $v_j \equiv 0$. \square

Example 3 (Complex Hopf fibrations). Consider the odd dimensional spheres \mathbb{S}^{2d+1}

$$\mathbb{S}^{2d+1} = \{(z_0, z_1, \dots, z_d) \in \mathbb{C}^{d+1} \mid \|z\| = 1\},$$

equipped with the standard round metric. The unit complex numbers $\mathbb{S}^1 = \{z \in \mathbb{C} \mid |z| = 1\}$ give an isometric action of $U(1)$ on \mathbb{S}^{2d+1} by

$$z \rightarrow e^{i\vartheta} z, \quad z \in \mathbb{S}^{2d+1}, \vartheta \in (-\pi, \pi].$$

Hence, the quotient space $\mathbb{S}^{2d+1}/\mathbb{S}^1 \simeq \mathbb{C}\mathbb{P}^d$ (the complex projective space) has a unique Riemannian structure (the Fubini-Study metric) such that the projection

$$p(z_0, \dots, z_d) = [z_0 : \dots : z_d]$$

is a Riemannian submersion. The fibration $\mathbb{S}^1 \hookrightarrow \mathbb{S}^{2d+1} \xrightarrow{p} \mathbb{C}\mathbb{P}^d$ is called the *complex Hopf fibration*. In real coordinates $z_j = x_j + iy_j$ on \mathbb{C}^{d+1} , the vertical distribution $\mathcal{V} = \ker p_*$ is generated by the restriction to \mathbb{S}^{2d+1} of the unit vector field

$$\xi = \sum_{j=0}^d (x_j \partial_{y_j} - y_j \partial_{x_j}).$$

The orthogonal complement $\mathcal{D} := \mathcal{V}^\perp$ with the restriction $g|_{\mathcal{D}}$ of the round metric define the standard sub-Riemannian structure on the complex Hopf fibrations. In real coordinates, as subspaces of \mathbb{R}^{2d+2} , the hemisphere and its boundary are

$$M = \mathbb{S}_+^{2d+1} := \left\{ \sum_{i=0}^d x_i^2 + y_i^2 = 1 \mid x_0 \geq 0 \right\}, \quad \partial M = \left\{ \sum_{i=0}^d x_i^2 + y_i^2 = 1 \mid x_0 = 0 \right\}.$$

A different set of coordinates we will use is the following

$$(\vartheta, w_1, \dots, w_d) \mapsto \left(\frac{e^{i\vartheta}}{\sqrt{1+|w|^2}}, \frac{w_1 e^{i\vartheta}}{\sqrt{1+|w|^2}}, \dots, \frac{w_d e^{i\vartheta}}{\sqrt{1+|w|^2}} \right),$$

where $\vartheta \in (-\pi, \pi)$ and $w = (w_1, \dots, w_d) \in \mathbb{C}^d$. In particular (w_1, \dots, w_d) are inhomogeneous coordinates for $\mathbb{C}\mathbb{P}^d$ given by $w_j = z_j/z_0$ and ϑ is the fiber coordinate. The north pole corresponds to $\vartheta = 0$ and $w = 0$. The hemisphere is characterized by $\vartheta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ and its boundary by $\cos(\vartheta) = 0$.

Example 4 (Quaternionic Hopf fibrations). Let \mathbb{H} be the field of quaternions. If $q = x + iy + jz + kw$, with $x, y, z, w \in \mathbb{R}$, the quaternionic norm is

$$\|q\| = x^2 + y^2 + z^2 + w^2.$$

Consider the sphere \mathbb{S}^{4d+3} as a subset of the quaternionic space \mathbb{H}^d ,

$$\mathbb{S}^{4d+3} = \{(q_0, q_1, \dots, q_d) \in \mathbb{H}^{d+1} \mid \|q\| = 1\},$$

equipped with the standard round metric. The left multiplication by unit quaternions $\mathbb{S}^3 = \{q \in \mathbb{H} \mid |q| = 1\}$ gives an isometric action of $\text{SU}(2)$ on \mathbb{S}^{4d+3} . The quotient space $\mathbb{S}^{4d+3}/\mathbb{S}^3 \simeq \mathbb{H}\mathbb{P}^d$ (the quaternionic projective space) has a unique Riemannian structure such that the projection

$$p(q_0, \dots, q_d) = [q_0 : \dots : q_d]$$

is a Riemannian submersion. The fibration $\mathbb{S}^3 \hookrightarrow \mathbb{S}^{4d+3} \xrightarrow{p} \mathbb{H}\mathbb{P}^d$ is the *quaternionic Hopf fibration*. In real coordinates $q_j = x_j + iy_j + jz_j + kw_j$ on \mathbb{H}^{d+1} , the vertical distribution $\mathcal{V} = \ker p_*$ is generated by

$$\xi_I = \sum_{i=0}^d y_i \partial_{x_i} - x_i \partial_{y_i} + w_i \partial_{z_i} - z_i \partial_{w_i}, \quad \xi_J = \sum_{i=0}^d z_i \partial_{x_i} - w_i \partial_{y_i} - x_i \partial_{z_i} + y_i \partial_{w_i},$$

$$\xi_K = \sum_{i=0}^d w_i \partial_{x_i} + z_i \partial_{y_i} - y_i \partial_{z_i} - x_i \partial_{w_i}.$$

The orthogonal complement $\mathcal{D} := \mathcal{V}^\perp$ with the restriction $g|_{\mathcal{D}}$ of the round metric define the standard sub-Riemannian structure on the quaternionic Hopf fibrations.

In real coordinates, the hemisphere $M = \mathbb{S}_+^{4d+4} \subset \mathbb{R}^{4d+4}$ and its boundary are

$$M = \left\{ \sum_{i=0}^d x_i^2 + y_i^2 + z_i^2 + w_i^2 = 1 \mid x_0 \geq 0 \right\},$$

$$\partial M = \left\{ \sum_{i=0}^d x_i^2 + y_i^2 + z_i^2 + w_i^2 = 1 \mid x_0 = 0 \right\}.$$

A different set of coordinates we will use is the following

$$(\vartheta_1, \vartheta_2, \vartheta_3, w_1, \dots, w_d) \mapsto \left(\frac{e^{i\vartheta_1 + j\vartheta_2 + k\vartheta_3}}{\sqrt{1+|w|^2}}, \frac{w_1 e^{i\vartheta_1 + j\vartheta_2 + k\vartheta_3}}{\sqrt{1+|w|^2}}, \dots, \frac{w_d e^{i\vartheta_1 + j\vartheta_2 + k\vartheta_3}}{\sqrt{1+|w|^2}} \right),$$

where $|\vartheta|^2 = \vartheta_1^2 + \vartheta_2^2 + \vartheta_3^2 < \pi^2$ and $w = (w_1, \dots, w_d) \in \mathbb{H}^d$. In particular (w_1, \dots, w_d) are inhomogeneous coordinates for $\mathbb{H}\mathbb{P}^d$ given by $w_j = q_0^{-1} q_j$ and $\vartheta_1, \vartheta_2, \vartheta_3$ are local

coordinates on $SU(2)$. The north pole corresponds to $\vartheta_1 = \vartheta_2 = \vartheta_3 = 0$ and $w = 0$. The hemisphere is characterized by $|\vartheta| \leq \pi/2$ and its boundary by $\cos|\vartheta| = 0$.

6. APPLICATIONS

In this section we present the proofs of the applications of the reduced Santaló formula presented in Sections 1.3, 1.4 and 1.5.

6.1. Hardy-type inequalities. It is well known that for all $f \in C_0^\infty([0, a])$ one has

$$(25) \quad \int_0^a f'(t)^2 dt \geq \frac{\pi^2}{a^2} \int_0^a f(t)^2 dt, \quad (1D \text{ Poincaré inequality})$$

with equality holding if and only if $f(t) = C \sin(\frac{\pi}{a}t)$. Moreover,

$$\int_0^a f'(t)^2 dt \geq \frac{1}{4} \int_0^a \frac{f(t)^2}{d(t)^2} dt, \quad (1D \text{ Hardy's inequality})$$

where $d(t) = \min\{t, a-t\}$ is the distance from the boundary and the equality holds if and only if $f(t) = 0$.

Recall that $\ell(\lambda)$ is the length at which the geodesic with initial covector λ leaves M crossing the boundary ∂M and, in general, $t \mapsto \ell(\phi_t(\lambda))$ is a decreasing function. Then $\ell(\cdot)$ is not invariant under the flow ϕ_t . For this reason in Section 1.3 we introduced the function $L : U^*M \rightarrow [0, +\infty]$ defined as $L(\lambda) := \ell(\lambda) + \ell(-\lambda)$, that measures the length of the projection of the maximal integral line of \vec{H} passing through λ . Indeed, $L(\cdot)$ is ϕ_t -invariant and coincides with $\ell(\cdot)$ on $U^+\partial M$, since it can be equivalently defined as

$$L(\lambda) := \begin{cases} \ell(-\phi_{\ell(-\lambda)}(-\lambda)), & \text{if } \ell(-\lambda) < +\infty, \\ +\infty, & \text{otherwise.} \end{cases}$$

Proof of Proposition 2. Choose coordinates x around a fixed $q \in M$ as in Lemma 18, and let (p, x) be the associated canonical coordinates on T^*M . Let Q be a quadratic form on $T_q^*M^r$. In particular $Q(\lambda) = \sum_{i,j=1}^k p_i Q_{ij} p_j$, where $\lambda = (p_1, \dots, p_k)$. By Lemma 18 we have

$$\int_{U_q^*M^r} Q(\lambda) \eta_{q_0}^r(\lambda) = \int_{\mathbb{S}^{k-1}} \sum_{i,j=1}^k Q_{ij} p_i p_j d\text{vol}_{\mathbb{S}^{k-1}} = \frac{|\mathbb{S}^{k-1}|}{k} \text{Trace}(Q),$$

where we performed the standard integral of a quadratic form on \mathbb{S}^{k-1} . Choosing $Q(\lambda) = \langle \lambda, \nabla_H f(q) \rangle^2$, then $\text{Trace}(Q) = |\nabla_H f(q)|^2$. Thus, for any point $q \in M$ we have

$$(26) \quad \frac{|\mathbb{S}^{k-1}|}{k} |\nabla_H f(q)|^2 = \int_{U_q^*M^r} \langle \lambda, \nabla_H f(q) \rangle^2 \eta_q^r(\lambda), \quad \forall f \in C^\infty(M).$$

Using the reduced Santaló formula (21),

$$\begin{aligned} \frac{|\mathbb{S}^{k-1}|}{k} \int_M |\nabla_H f(q)|^2 \omega(q) &= \int_M \left[\int_{U_q^*M^r} \langle \lambda, \nabla_H f(q) \rangle^2 \eta_q^r(\lambda) \right] \omega(q) \\ &= \int_{U^*M^r} \langle \lambda, \nabla_H f \rangle^2 \mu^r \\ &\geq \int_{U^{\circlearrowleft} M^r} \langle \lambda, \nabla_H f \rangle^2 \mu^r \\ &= \int_{\partial M} \left[\int_{U_q^+ \partial M^r} \left(\int_0^{\ell(\lambda)} \langle \phi_t(\lambda), \nabla_H f \rangle^2 dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q^r(\lambda) \right] \sigma(q). \end{aligned}$$

Consider the subset $D = U^+ \partial M^r \cap \{\ell < +\infty\}$. Let $f_\lambda(t) := f(\pi \circ \phi_t(\lambda))$. For $\lambda \in D$ we have $f_\lambda(0) = f_\lambda(\ell(\lambda)) = 0$ and the one-dimensional Poincaré inequality (25) gives

$$(27) \quad \int_0^{\ell(\lambda)} \langle \phi_t(\lambda), \nabla_H f \rangle^2 dt = \int_0^{\ell(\lambda)} f'_\lambda(t)^2 dt \geq \frac{\pi^2}{\ell^2(\lambda)} \int_0^{\ell(\lambda)} f_\lambda(t)^2 dt.$$

Indeed we can replace ℓ with L , which is ϕ_t -invariant. Then

$$\frac{|\mathbb{S}^{k-1}|}{k} \int_M |\nabla_H f(q)|^2 \omega(q) \geq \pi^2 \int_{\partial M} \left[\int_{D_q} \left(\int_0^{\ell(\lambda)} \frac{f_\lambda(t)^2}{L(\lambda)^2} dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q^r(\lambda) \right] \sigma(q).$$

Since on $U_q^+ \partial M \setminus D_q$ the function $1/L(\lambda)^2 = 0$, we can replace D_q with $U_q^+ \partial M$. Using again Santaló formula to restore the integral on $U_q^* M^r$, we obtain

$$\int_M |\nabla_H f(q)|^2 \omega(q) \geq \frac{k\pi^2}{|\mathbb{S}^{k-1}|} \int_{U_q^* M^r} \frac{(\pi^* f)^2}{L^2} \mu^r = \frac{k\pi^2}{|\mathbb{S}^{k-1}|} \int_M \left[\int_{U_q^* M^r} \frac{1}{L^2} \eta_q^r \right] f(q)^2 \omega(q).$$

The second equality follows by Lemma 11. This concludes the proof of (4).

To prove (5) we replace Poincaré inequality with Hardy's in (27):

$$\begin{aligned} \int_0^{\ell(\lambda)} \langle \lambda, \nabla_H f \rangle^2 dt &= \int_0^{\ell(\lambda)} f'_\lambda(t)^2 dt \\ &\geq \frac{1}{4} \int_0^{\ell(\lambda)} \frac{f_\lambda(t)^2}{\min\{t, \ell(\lambda) - t\}^2} dt \geq \frac{1}{4} \int_0^{\ell(\lambda)} \frac{f(\phi_t(\lambda(t)))^2}{\ell(\phi_t(\lambda))^2} dt. \end{aligned}$$

We then proceed as in the previous case without replacing ℓ with L . \square

Proof of Proposition 3. The result is obtained by mimicking the proof of Proposition 2. Observe that, for any $f \in C_0^\infty(M)$ and $q \in M$, we have

$$\begin{aligned} \int_{U_q^* M^r} |\langle \lambda, \nabla_H f(q) \rangle|^p \eta_q^r(\lambda) &= |\nabla_H f(q)|^p \int_{U_q^* M^r} \left| \langle \lambda, \frac{\nabla_H f(q)}{|\nabla_H f(q)|} \rangle \right|^p \eta_q^r(\lambda) \\ &= 2 |\nabla_H f(q)|^p \int_{\mathbb{S}^{k-1} \cap \{p_1 > 0\}} p_1^p \text{vol}_{\mathbb{S}^{k-1}} \\ &= |\nabla_H f(q)|^p C_{p,k}, \end{aligned}$$

where we used coordinates as in Lemma 18, the rotational invariance of the measure and

$$C_{p,k} = \frac{k}{|\mathbb{S}^{k-1}|} \frac{2\Gamma(1 + \frac{k}{2})\Gamma(1 + \frac{p}{2})}{\sqrt{\pi}\Gamma(\frac{k+p}{2})}.$$

Using the above in place of (26), by Santaló formula (21) we obtain

$$\frac{|\mathbb{S}^{k-1}|}{k} C_{p,k} \int_M |\nabla_H f|^p \omega(q) \geq \int_{\partial M} \left[\int_{U_q^+ \partial M^r} \left(\int_0^{\ell(\lambda)} |\langle \phi_t(\lambda), \nabla_H f \rangle|^p dt \right) \langle \lambda, \mathbf{n}_q \rangle \eta_q^r(\lambda) \right] \sigma(q).$$

To prove (6) we proceed as in the proof of Proposition 2 replacing the step (27) with the L^p Poincaré inequality [37, Ch. 3]

$$\int_0^a |f'(t)|^p dt \geq \left(\frac{\pi_p}{a} \right)^p \int_0^a |f(t)|^p dt.$$

We proceed similarly for the proof of (7), replacing (27) with the L^p Hardy's inequalities [31, Thm. 327]

$$\int_0^a |f'(t)|^p dt \geq \left(\frac{p-1}{p} \right)^p \int_0^a \frac{|f(t)|^p}{d(t)^p} dt. \quad \square$$

Proof of Proposition 4. With $L := \sup_{\lambda \in U^*M^r} L(\lambda)$, the Hardy inequality (4) can be further simplified into

$$\int_M |\nabla_H f|^2 \omega \geq \frac{k\pi^2}{L^2} \int_M f^2 \omega.$$

By the min-max principle (13), whenever any $f \in C_0^\infty(M)$ such that $\int_M f^2 \omega = 1$, we have

$$\lambda_1(M) \geq \int_M |\nabla_H f|^2 \omega \geq \frac{k\pi^2}{L^2}. \quad \square$$

Proof of Proposition 5. Fix a north pole q_0 and the hemisphere M whose center is q_0 . By Remark 8, in all three cases, the reduced sub-Riemannian geodesics are a subset of the Riemannian ones (great circles). In particular $L = \pi$ and Proposition 4 gives

$$\lambda_1(M) \geq k,$$

where $k = d$ for the Riemannian sphere, $k = 2d$ for the CHF and $k = 4d$ for the QHF.

In all cases, uniqueness of $\Phi = \cos(\delta) \in C_0^\infty(M)$ follows as in the Riemannian case from the min-max principle [18, Corollary 2, p. 20]. To complete the proof, we show that the function Φ is an eigenfunction of the positive (sub-)Laplacian with eigenvalue $d, 2d, 4d$, respectively. In the Riemannian case this is well known. For the CHF we use coordinates (ϑ, w) of Example 3. Then,

$$\Phi = x_0 = \frac{\cos(\vartheta)}{\sqrt{1+|w|^2}} = \cos(\vartheta) \cos(r),$$

where we have set $\tan(r) = |w|$. In [8, Proposition 2.3] the authors show that for a function depending only on ϑ and r the action of the sub-Laplacian reduces to the action of its cylindrical part, given by

$$\tilde{\Delta} = \partial_r^2 + ((2d-1)\cot(r) - \tan(r))\partial_r + \tan^2(r)\partial_\vartheta^2.$$

In particular $\Delta\Phi = \tilde{\Delta}\Phi = (-2d)\Phi$.

For the QHF we use coordinates $(\vartheta_1, \vartheta_2, \vartheta_3, w)$ of Example 4. Using the expression

$$e^{i\vartheta_1 + j\vartheta_2 + k\vartheta_3} = \cos(\eta) + (i\vartheta_1 + j\vartheta_2 + k\vartheta_3) \frac{\sin(\eta)}{\eta}, \quad \eta := \sqrt{\vartheta_1^2 + \vartheta_2^2 + \vartheta_3^2},$$

we obtain

$$\Phi = x_0 = \frac{\cos(\eta)}{\sqrt{1+|w|^2}} = \cos(\eta) \cos(r),$$

where we have set $\tan(r) = |w|$. In [9, Definition 2.1 and Proposition 2.2] the authors show that, for a function depending only on η and r , the action of the sub-Laplacian reduces to the action of its cylindrical part, given by

$$\tilde{\Delta} = \partial_r^2 + ((4d-1)\cot(r) - 3\tan(r))\partial_r + \tan^2(r) \left(\partial_\eta^2 + 2\cot(\eta)\partial_\eta \right).$$

In particular $\Delta\Phi = \tilde{\Delta}\Phi = (-4d)\Phi$. □

6.2. Isoperimetric inequalities. We define some quantities that we already introduced.

Definition 6. The *visibility angle* at $q \in M$ and the *optimal visibility angle* are

$$\vartheta_q^\star := \frac{\eta_q^r(U_q^\star M^r)}{\eta_q^r(U_q M^r)}, \quad \tilde{\vartheta}_q^\star := \frac{\eta_q^r(\tilde{U}_q^\star M^r)}{\eta_q^r(U_q M^r)}.$$

The *least visibility angle* and the *least optimal visibility angle* are

$$\vartheta^\star = \inf_{q \in M} \vartheta_q^\star, \quad \tilde{\vartheta}^\star = \inf_{q \in M} \tilde{\vartheta}_q^\star.$$

Notice that $\vartheta_q^\star, \tilde{\vartheta}_q^\star, \vartheta^\star, \tilde{\vartheta}^\star \in [0, 1]$ and do not depend on the choice of the volume ω .

Definition 7. The sub-Riemannian *diameter* and *reduced diameter* are:

$$\begin{aligned} \text{diam}(M) &:= \sup\{\mathbf{d}(x, y) \mid x, y \in M\} = \sup\{\tilde{\ell}(\lambda) \mid \lambda \in U^*M\}, \\ \text{diam}^r(M) &:= \sup\{\tilde{\ell}(\lambda) \mid \lambda \in U^*M^r\}. \end{aligned}$$

Clearly $\text{diam}^r(M) \leq \text{diam}(M)$.

Proof of Proposition 7. The proof follows as in [19, 21], considering $F = 1$ in (21). The l.h.s. is estimated from below using the disintegration of μ^r given in Lemma 11. For the estimate of the r.h.s. we only observe that, by Lemma 18 we have

$$\int_{U_q^+ \partial M^r} \langle \lambda, \mathbf{n}_q \rangle \eta_q^r(\lambda) = \int_{\mathbb{S}^{k-1} \cap \{p_1 > 0\}} p_1 \text{vol}_{\mathbb{S}^{k-1}} = \frac{|\mathbb{S}^{k-2}|}{k-1} = \frac{|\mathbb{S}^k|}{2\pi}. \quad \square$$

Proof of Proposition 8. For all these structures, all the inequalities in the proof of Proposition 7 are equalities, hence the sharpness follows. Anyway, here we perform the explicit computation for the hemisphere of the sub-Riemannian complex Hopf fibration; the remaining case of the quaternionic Hopf fibration can be checked following the same steps.

We use the notation of Example 3, and real coordinates. Let $q = (0, y_0, \dots, x_d, y_d) \in \partial M$. The sub-Riemannian normal \mathbf{n}_q is the unique inward pointing unit vector in \mathcal{D}_q orthogonal to $\mathcal{D}_q \cap T_q \partial M$. Indeed, $T_q \partial M$ is the orthogonal complement to ∂_{x_0} w.r.t. to the Riemannian round metric, while \mathcal{D}_q is the orthogonal complement to ξ . Thus, $\mathbf{n}_q = \alpha \xi + \beta \partial_{x_0}$. The condition $\mathbf{n}_q \in \mathcal{D}$ and the normalization imply

$$\mathbf{n}_q = \frac{1}{\sqrt{1 - g(\partial_{x_0}, \xi)^2}} (\partial_{x_0} - g(\partial_{x_0}, \xi) \xi).$$

Using the explicit expression for ξ we obtain

$$\mathbf{n}_q = \sqrt{1 - y_0^2} \partial_{x_0} \quad \text{mod } T_q \partial M.$$

Notice that $C(\partial M) = \{y_0^2 = 1\} \cap \partial M$. Due to the factor $\rho(y_0) := \sqrt{1 - y_0^2}$, the sub-Riemannian surface measure σ is different from the Riemannian one in cylindrical coordinates $\sigma_R = \iota_{\partial_{x_0}} \omega = \rho^{2d-2} dy_0 d\mu_{\mathbb{S}^{2d-1}}$. In particular

$$\sigma(\partial M) = \int_{\partial M} \rho(y_0) \sigma_R = |\mathbb{S}^{2d-2}| \int_{-1}^1 \rho(y_0)^{2d-1} dy_0 = \frac{|\mathbb{S}^{2d+1}|}{|\mathbb{S}^{2d}|}.$$

Moreover $\omega(M) = |\mathbb{S}^{2d+1}|/2$. By Remark 8, the reduced geodesics γ_λ with $\lambda \in U^*M^r$ are a subset of Riemannian geodesics hence $\vartheta^{\check{\bullet}} = \check{\vartheta}^{\check{\bullet}} = 1$ and $\ell = \text{diam}^r(M) = \pi$. \square

APPENDIX A. LIE DERIVATIVE OF THE REDUCED LIOUVILLE VOLUME

Lemma 24. *In the notation of Section 4.3, we have*

$$\mathcal{L}_{\vec{H}} \Theta^r = - \left(\sum_{j=1}^{n-k} \sum_{i=1}^k u_i \partial_{v_j} \{u_i, v_j\} \right) \Theta^r.$$

Proof. For any ℓ -uple $w = (w_1, \dots, w_\ell)$ of vector fields and ℓ -form α , we denote

$$\alpha(\mathcal{L}_{\vec{H}}(w)) = \sum_{i=1}^{\ell} \alpha(w_1, \dots, [\vec{H}, w_i], \dots, w_\ell).$$

Let $(x_1, \dots, x_n) : U \rightarrow \mathbb{R}^n$ be coordinates on $U \subset M$. Then $(x, u, v) : \pi^{-1}(U) \rightarrow \mathbb{R}^{2n}$ are local coordinates for T^*M and $T^*M^r \cap \pi^{-1}(U) = \{(x, u, v) \mid v = 0\}$.

Denote $\partial_v = (\partial_{v_1}, \dots, \partial_{v_{n-k}})$ and $\partial_u = (\partial_{u_1}, \dots, \partial_{u_k})$ and $\partial_x = (\partial_{x_1}, \dots, \partial_{x_n})$. Recall that $\Theta^r(\partial_u, \partial_x) = \Theta(\partial_u, \partial_v, \partial_x) = (-1)^{k(n-k)}\Theta(\partial_v, \partial_u, \partial_x)$. Then, using twice $(\mathcal{L}_{\vec{H}}\alpha)(w) = \vec{H}(\alpha(w)) - \alpha(\mathcal{L}_{\vec{H}}(w))$ for any ℓ -form α and ℓ -uple w , we obtain

$$\begin{aligned}
 (\mathcal{L}_{\vec{H}}\Theta^r)(\partial_u, \partial_x) &= \vec{H}(\Theta^r(\partial_u, \partial_x)) - \Theta^r(\mathcal{L}_{\vec{H}}(\partial_u, \partial_x)) \\
 &= (-1)^{k(n-k)} \left[\vec{H}(\Theta(\partial_v, \partial_u, \partial_x)) - \Theta(\partial_v, \mathcal{L}_{\vec{H}}(\partial_u, \partial_x)) \right] \\
 &= (-1)^{k(n-k)} \left[(\mathcal{L}_{\vec{H}}\Theta)(\partial_v, \partial_u, \partial_x) + \Theta(\mathcal{L}_{\vec{H}}(\partial_v), \partial_u, \partial_x) \right] \\
 (28) \qquad \qquad \qquad &= \Theta(\partial_u, \mathcal{L}_{\vec{H}}(\partial_v), \partial_x),
 \end{aligned}$$

where, in the last step, we used that $\mathcal{L}_{\vec{H}}\Theta = 0$. Now observe that, for $j = 1, \dots, n-k$,

$$\begin{aligned}
 (\vec{H}, \partial_{v_j}) &= \sum_{i=1}^k [u_i \vec{u}_i, \partial_{v_j}] = \sum_{i=1}^k u_i [\vec{u}_i, \partial_{v_j}] \\
 (29) \qquad \qquad \qquad &= - \sum_{i=1}^k \sum_{\ell=1}^k u_i \partial_{v_j} \{u_i, u_\ell\} \partial_{u_\ell} - \sum_{i=1}^k \sum_{\ell=1}^{n-k} u_i \partial_{v_j} \{u_i, v_\ell\} \partial_{v_\ell}.
 \end{aligned}$$

Plugging (29) in (28), and using complete skew-symmetry, we obtain the statement. \square

ACKNOWLEDGMENTS

We are grateful to the organizers of the Trimester *Geometry, Analysis and Dynamics on Sub-Riemannian Manifolds*, whose stimulating atmosphere fostered this collaboration, and the Institut Henri Poincaré, Paris, where most of this research has been carried out. We also thank the Isaac Newton Institute for Mathematical Sciences, Cambridge, for the support and hospitality during the programme *Periodic and Ergodic Spectral Problems*. The last named author is grateful for the support and hospitality of the Erwin Schrödinger Institute of Mathematical Physics in Vienna during the programme *Modern theory of wave equations*, where he stayed when this article was completed. We thank F. Baudoin and J. Wang for useful discussions about the spectrum of the sub-Laplacian on the quaternionic Hopf fibration and L. Parnowski for useful comments on the spectral theoretic part.

REFERENCES

- [1] A. Agrachev. Some open problems. In *Geometric control theory and sub-Riemannian geometry*, volume 5 of *Springer INdAM Ser.*, pages 1–13. Springer, Cham, 2014.
- [2] A. Agrachev, D. Barilari, and U. Boscain. *Introduction to Riemannian and Sub-Riemannian geometry from Hamiltonian viewpoint (Lecture notes)*. 2015. <http://www.math.jussieu.fr/~barilari/Notes.php>.
- [3] G. Barbatis and A. Tertikas. On the Hardy constant of some non-convex planar domains. *ArXiv e-prints*, Sept. 2014.
- [4] D. Barilari and L. Rizzi. A formula for Popp’s volume in sub-Riemannian geometry. *Anal. Geom. Metr. Spaces*, 1:42–57, 2013.
- [5] F. Baudoin. *Sub-Laplacians and hypoelliptic operators on totally geodesic Riemannian foliations*. Oct. 2014. Insitute Henri Poincaré course.
- [6] F. Baudoin and M. Bonnefont. The subelliptic heat kernel on $SU(2)$: representations, asymptotics and gradient bounds. *Math. Z.*, 263(3):647–672, 2009.
- [7] F. Baudoin and B. Kim. Sobolev, Poincaré, and isoperimetric inequalities for subelliptic diffusion operators satisfying a generalized curvature dimension inequality. *Revista Matemática Iberoamericana*, 30(1):109–131, 2014.
- [8] F. Baudoin and J. Wang. The subelliptic heat kernel on the CR sphere. *Math. Z.*, 275(1-2):135–150, 2013.
- [9] F. Baudoin and J. Wang. The subelliptic heat kernels of the quaternionic Hopf fibration. *Potential Anal.*, 41(3):959–982, 2014.
- [10] A. L. Besse. *Einstein manifolds*. Classics in Mathematics. Springer-Verlag, Berlin, 2008. Reprint of the 1987 edition.
- [11] U. Boscain and C. Laurent. The Laplace-Beltrami operator in almost-Riemannian geometry. *Ann. Inst. Fourier (Grenoble)*, 63(5):1739–1770, 2013.

- [12] U. Boscain, R. Neel, and L. Rizzi. Intrinsic random walks and sub-Laplacians in sub-Riemannian geometry. *ArXiv e-prints*, Mar. 2015.
- [13] U. Boscain, D. Prandi, and M. Seri. Spectral analysis and the Aharonov-Bohm effect on certain almost-Riemannian manifolds. *Comm. in PDE (to appear)*, 2015.
- [14] H. Brezis and M. Marcus. Hardy's inequalities revisited. *Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4)*, 25(1-2):217–237 (1998). Dedicated to Ennio De Giorgi.
- [15] L. Capogna, D. Danielli, and N. Garofalo. An isoperimetric inequality and the geometric Sobolev embedding for vector fields. *Math. Res. Lett.*, 1(2):263–268, 1994.
- [16] L. Capogna, D. Danielli, S. D. Pauls, and J. T. Tyson. *An introduction to the Heisenberg group and the sub-Riemannian isoperimetric problem*, volume 259 of *Progress in Mathematics*. Birkhäuser Verlag, Basel, 2007.
- [17] S. Chanillo and P. C. Yang. Isoperimetric inequalities & volume comparison theorems on CR manifolds. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 8(2):279–307, 2009.
- [18] I. Chavel. *Eigenvalues in Riemannian geometry*, volume 115 of *Pure and Applied Mathematics*. Academic Press, Inc., Orlando, FL, 1984. Including a chapter by Burton Randol, With an appendix by Jozef Dodziuk.
- [19] I. Chavel. *Riemannian geometry*, volume 98 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2006.
- [20] Y. Colin de Verdière, L. Hillairet, and E. Trélat. Spectral asymptotics for sub-Riemannian Laplacians. I: quantum ergodicity and quantum limits in the 3D contact case. *ArXiv e-prints*, Apr. 2015.
- [21] C. B. Croke. Some isoperimetric inequalities and eigenvalue estimates. *Ann. Sci. École Norm. Sup. (4)*, 13(4):419–435, 1980.
- [22] C. B. Croke. A sharp four-dimensional isoperimetric inequality. *Comment. Math. Helv.*, 59(2):187–192, 1984.
- [23] C. B. Croke and A. Derdziński. A lower bound for λ_1 on manifolds with boundary. *Comment. Math. Helv.*, 62(1):106–121, 1987.
- [24] D. Danielli, N. Garofalo, and N. C. Phuc. Hardy-Sobolev Type Inequalities with Sharp Constants in Carnot-Carathéodory Spaces. *Potential Analysis*, 34(3):223–242, 2011.
- [25] E. B. Davies. A review of Hardy inequalities. In *The Mał'ya anniversary collection, Vol. 2 (Rostock, 1998)*, volume 110 of *Oper. Theory Adv. Appl.*, pages 55–67. Birkhäuser, Basel, 1999.
- [26] T. Ekholm, H. Kovařík, and A. Laptev. Hardy inequalities for p -Laplacians with Robin boundary conditions. *Nonlinear Anal.*, 128:365–379, 2015.
- [27] R. H. Escobales, Jr. Riemannian submersions with totally geodesic fibers. *J. Differential Geom.*, 10:253–276, 1975.
- [28] B. Green and T. Tao. Linear equations in primes. *Ann. of Math. (2)*, 171(3):1753–1850, 2010.
- [29] M. Gromov. Almost flat manifolds. *J. Differential Geom.*, 13(2):231–241, 1978.
- [30] A. M. Hansson and A. Laptev. Sharp spectral inequalities for the Heisenberg Laplacian. In *Groups and analysis*, volume 354 of *London Math. Soc. Lecture Note Ser.*, pages 100–115. Cambridge Univ. Press, Cambridge, 2008.
- [31] G. H. Hardy, J. E. Littlewood, and G. Pólya. *Inequalities*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1988. Reprint of the 1952 edition.
- [32] L. Hörmander. Hypoelliptic second order differential equations. *Acta Math.*, 119:147–171, 1967.
- [33] S. Ivanov and D. Vassilev. The Lichnerowicz and Obata first eigenvalue theorems and the Obata uniqueness result in the Yamabe problem on CR and quaternionic contact manifolds. *Nonlinear Analysis: Theory, Methods & Applications*, 126:262 – 323, 2015.
- [34] E. Le Donne, G. P. Leonardi, R. Monti, and D. Vittone. Extremal curves in nilpotent Lie groups. *Geom. Funct. Anal.*, 23(4):1371–1401, 2013.
- [35] E. Le Donne, R. Montgomery, A. Ottazzi, P. Pansu, and D. Vittone. Sard Property for the endpoint map on some Carnot groups. *Annales de l'Institut Henri Poincaré, Analyse non linéaire (to appear)*, Mar. 2015.
- [36] J. M. Lee. *Introduction to smooth manifolds*, volume 218 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2013.
- [37] P. Lindqvist. *A nonlinear eigenvalue problem (Lecture Notes)*. 2000. <http://www.math.ntnu.no/~lqvist/nonlineigen.pdf>.
- [38] F. Montefalcone. Some relations among volume, intrinsic perimeter and one-dimensional restrictions of BV functions in Carnot groups. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 4(1):79–128, 2005.
- [39] R. Montgomery. *A tour of subriemannian geometries, their geodesics and applications*, volume 91 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2002.
- [40] P. Pansu. An isoperimetric inequality on the Heisenberg group. *Rend. Sem. Mat. Univ. Politec. Torino*, (Special Issue):159–174 (1984), 1983. Conference on differential geometry on homogeneous spaces (Turin, 1983).

- [41] L. Rifford. *Sub-Riemannian Geometry and Optimal Transport*. SpringerBriefs in Mathematics. Springer, 2014.
- [42] L. Rizzi and P. Silveira. Sub-Riemannian Ricci curvatures and universal diameter bounds for 3-Sasakian structures. *ArXiv e-prints*.
- [43] L. A. Santaló. *Integral geometry and geometric probability*. Cambridge Mathematical Library. Cambridge University Press, Cambridge, second edition, 2004. With a foreword by Mark Kac.
- [44] R. S. Strichartz. Estimates for sums of eigenvalues for domains in homogeneous spaces. *J. Funct. Anal.*, 137(1):152–190, 1996.
- [45] L. Yuan and W. Zhao. Some formulas of Santaló type in Finsler geometry and its applications. *ArXiv e-prints*, Apr. 2014.